



# CANADIAN STROKE BEST PRACTICE RECOMMENDATIONS

**Rehabilitation, Recovery and Community Participation  
Following Stroke**

**Part Two: Delivery of Stroke Rehabilitation to Optimize  
Functional Recovery**

**7<sup>th</sup> Edition, 2025**

## **Appendix Three: References**

*Salbach N, Yao J (Writing Group Co-Chairs)  
on behalf of the Canadian Stroke Best Practice Recommendations  
Stroke Rehabilitation and Recovery Writing Group*

© 2025 Heart and Stroke Foundation of Canada

## APPENDIX THREE: REFERENCES

1. Nelson MLA, Hanna E, Hall S, Calvert M. What makes stroke rehabilitation patients complex? Clinician perspectives and the role of discharge pressure. *J Comorb.* 2016;6:35-41.
2. Holodinsky JK, Lindsay P, Yu AYX, Ganesh A, Joundi RA, Hill MD. Estimating the Number of Hospital or Emergency Department Presentations for Stroke in Canada. *Can J Neurol Sci.* 2023;50:820-825.
3. Government of Canada. Canadian Chronic Disease Surveillance System (CCDSS). 2023; <https://health-infobase.canada.ca/ccdss/data-tool/Index>. Accessed March 3, 2025.
4. Kapral MK, Hall R, Fang J, et al. Predictors of Hospitalization in Patients With Transient Ischemic Attack or Minor Ischemic Stroke. *Can J Neurol Sci.* 2016;43:523-528.
5. Canadian Institutes of Health Research. Transitions in Care: Overview. 2019; <http://www.cihr-irsc.gc.ca/e/50972.html> Accessed October 31, 2024.
6. World Health Organization. International Classification of Functioning, Disability and Health. 2001; <https://iris.who.int/bitstream/handle/10665/42407/9241545429-eng.pdf>. Accessed June 12, 2024.
7. Li S, Francisco GE, Rymer WZ. A New Definition of Poststroke Spasticity and the Interference of Spasticity With Motor Recovery From Acute to Chronic Stages. *Neurorehabil Neural Repair.* 2021;35:601-610.
8. Canadian Stroke Best Practices. Canadian Stroke Best Practice Recommendations. Overview of Methodology 7<sup>th</sup> Edition 2019-2023; <https://www.strokebestpractices.ca/recommendations/overview-methods-and-knowledge-translation>. Accessed March 3, 2025.
9. Brouwers MC, Kho ME, Browman GP, et al. AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ.* 2010;182:E839-842.
10. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol.* 2011;64:383-394.
11. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ.* 2008;336:924-926.
12. Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schünemann HJ. What is "quality of evidence" and why is it important to clinicians? *BMJ.* 2008;336:995-998.
13. Schünemann H BJ, Guyatt G, Oxman A, editors. GRADE handbook for grading quality of evidence and strength of recommendations. 2013; <https://guidelinedevelopment.org/handbook>. Accessed March 3, 2025.
14. French B, Thomas LH, Coupe J, et al. Repetitive task training for improving functional ability after stroke. *Cochrane Database Syst Rev.* 2016;11:Cd006073.

15. Langhorne P, Coupar F, Pollock A. Motor recovery after stroke: a systematic review. *The Lancet. Neurology*. 2009;8:741-754.
16. Shimodozono M, Noma T, Nomoto Y, et al. Benefits of a repetitive facilitative exercise program for the upper paretic extremity after subacute stroke: a randomized controlled trial. *Neurorehabil Neural Repair*. 2013;27:296-305.
17. Tenberg S, Mueller S, Vogt L, et al. Comparative Effectiveness of Upper Limb Exercise Interventions in Individuals With Stroke: A Network Meta-Analysis. *Stroke*. 2023;54:1839-1853.
18. Khan MA, Fares H, Ghayvat H, et al. A systematic review on functional electrical stimulation based rehabilitation systems for upper limb post-stroke recovery. *Front Neurol*. 2023;14:1272992.
19. Eraifej J, Clark W, France B, Desando S, Moore D. Effectiveness of upper limb functional electrical stimulation after stroke for the improvement of activities of daily living and motor function: a systematic review and meta-analysis. *Syst Rev*. 2017;6:40.
20. Vafadar AK, Côté JN, Archambault PS. Effectiveness of functional electrical stimulation in improving clinical outcomes in the upper arm following stroke: a systematic review and meta-analysis. *Biomed Res Int*. 2015;2015:729768.
21. Dromerick AW, Lang CE, Birkenmeier RL, et al. Very Early Constraint-Induced Movement during Stroke Rehabilitation (VECTORS): A single-center RCT. *Neurology*. 2009;73:195-201.
22. Wolf SL, Winstein CJ, Miller JP, et al. Effect of constraint-induced movement therapy on upper extremity function 3 to 9 months after stroke: the EXCITE randomized clinical trial. *JAMA*. 2006;296:2095-2104.
23. Page SJ, Boe S, Levine P. What are the "ingredients" of modified constraint-induced therapy? An evidence-based review, recipe, and recommendations. *Restor Neurol Neurosci*. 2013;31:299-309.
24. Kwakkel G, Winters C, van Wegen EE, et al. Effects of Unilateral Upper Limb Training in Two Distinct Prognostic Groups Early After Stroke: The EXPLICIT-Stroke Randomized Clinical Trial. *Neurorehabil Neural Repair*. 2016;30:804-816.
25. Liu XH, Huai J, Gao J, Zhang Y, Yue SW. Constraint-induced movement therapy in treatment of acute and sub-acute stroke: a meta-analysis of 16 randomized controlled trials. *Neural Regen Res*. 2017;12:1443-1450.
26. Corbetta D, Sirtori V, Castellini G, Moja L, Gatti R. Constraint-induced movement therapy for upper extremities in people with stroke. *Cochrane Database Syst Rev*. 2015;2015:Cd004433.
27. Gao Q, Zhang Y, Long J, Pan M, Wang J, Yang F. Effect of different constraint-induced movement therapy protocols on recovery of stroke survivors with upper extremity dysfunction: a systematic review and network meta-analysis. *Int J Rehabil Res*. 2023;46:133-150.
28. Thieme H, Morkisch N, Mehrholz J, et al. Mirror therapy for improving motor function after stroke. *Cochrane Database Syst Rev*. 2018;7:Cd008449.

29. Yang Y, Zhao Q, Zhang Y, Wu Q, Jiang X, Cheng G. Effect of Mirror Therapy on Recovery of Stroke Survivors: A Systematic Review and Network Meta-analysis. *Neuroscience*. 2018;390:318-336.
30. Zeng W, Guo Y, Wu G, Liu X, Fang Q. Mirror therapy for motor function of the upper extremity in patients with stroke: A meta-analysis. *J Rehabil Med*. 2018;50:8-15.
31. Barclay RE, Stevenson TJ, Poluha W, Semenko B, Schubert J. Mental practice for treating upper extremity deficits in individuals with hemiparesis after stroke. *Cochrane Database Syst Rev*. 2020;5:Cd005950.
32. Chen J, Or CK, Chen T. Effectiveness of Using Virtual Reality-Supported Exercise Therapy for Upper Extremity Motor Rehabilitation in Patients With Stroke: Systematic Review and Meta-analysis of Randomized Controlled Trials. *J Med Internet Res*. 2022;24:e24111.
33. Laver KE, Lange B, George S, Deutsch JE, Saposnik G, Crotty M. Virtual reality for stroke rehabilitation. *Cochrane Database Syst Rev*. 2017;11:Cd008349.
34. Adie K, Schofield C, Berrow M, et al. Does the use of Nintendo Wii Sports(TM) improve arm function? Trial of Wii(TM) in Stroke: a randomized controlled trial and economics analysis. *Clin Rehabil*. 2017;31:173-185.
35. Brunner I, Skouen JS, Hofstad H, et al. Virtual Reality Training for Upper Extremity in Subacute Stroke (VIRTUES): A multicenter RCT. *Neurology*. 2017;89:2413-2421.
36. Saposnik G, Cohen LG, Mamdani M, et al. Efficacy and safety of non-immersive virtual reality exercising in stroke rehabilitation (EVREST): a randomised, multicentre, single-blind, controlled trial. *The Lancet. Neurology*. 2016;15:1019-1027.
37. Kong KH, Loh YJ, Thia E, et al. Efficacy of a Virtual Reality Commercial Gaming Device in Upper Limb Recovery after Stroke: A Randomized, Controlled Study. *Top Stroke Rehabil*. 2016;23:333-340.
38. Harris JE, Eng JJ, Miller WC, Dawson AS. A self-administered Graded Repetitive Arm Supplementary Program (GRASP) improves arm function during inpatient stroke rehabilitation: a multi-site randomized controlled trial. *Stroke*. 2009;40:2123-2128.
39. Hunter SM, Johansen-Berg H, Ward N, et al. Functional Strength Training and Movement Performance Therapy for Upper Limb Recovery Early Poststroke-Efficacy, Neural Correlates, Predictive Markers, and Cost-Effectiveness: FAST-INDiCATE Trial. *Front Neurol*. 2017;8:733.
40. Harris JE, Eng JJ. Strength training improves upper-limb function in individuals with stroke: a meta-analysis. *Stroke*. 2010;41:136-140.
41. Gnanaprakasam A, Karthikbabu S, Ravishankar N, Solomon JM. Effect of task-based bilateral arm training on upper limb recovery after stroke: A systematic review and meta-analysis. *J Stroke Cerebrovasc Dis*. 2023;32:107131.

42. Chen S, Qiu Y, Bassile CC, Lee A, Chen R, Xu D. Effectiveness and Success Factors of Bilateral Arm Training After Stroke: A Systematic Review and Meta-Analysis. *Front Aging Neurosci.* 2022;14:875794.
43. van Delden AE, Peper CE, Beek PJ, Kwakkel G. Unilateral versus bilateral upper limb exercise therapy after stroke: a systematic review. *J Rehabil Med.* 2012;44:106-117.
44. Coupar F, Pollock A, van Wijck F, Morris J, Langhorne P. Simultaneous bilateral training for improving arm function after stroke. *Cochrane Database Syst Rev.* 2010;2010:Cd006432.
45. Thijs L, Voets E, Denissen S, et al. Trunk training following stroke. *Cochrane Database Syst Rev.* 2023;3:Cd013712.
46. Zhang Q, Fu C, Liang Z, et al. The effect of adding trunk restraint to task-oriented training in improving function in stroke patients: A systematic review and meta-analysis. *NeuroRehabilitation.* 2020;46:95-108.
47. Alexander J, Dawson J, Langhorne P. Dynamic hand orthoses for the recovery of hand and arm function in adults after stroke: A systematic review and meta-analysis of randomised controlled trials. *Top Stroke Rehabil.* 2022;29:114-124.
48. Elsner B, Kugler J, Pohl M, Mehrholz J. Transcranial direct current stimulation (tDCS) for improving activities of daily living, and physical and cognitive functioning, in people after stroke. *Cochrane Database Syst Rev.* 2020;11:Cd009645.
49. Chhatbar PY, Ramakrishnan V, Kautz S, George MS, Adams RJ, Feng W. Transcranial Direct Current Stimulation Post-Stroke Upper Extremity Motor Recovery Studies Exhibit a Dose-Response Relationship. *Brain Stimul.* 2016;9:16-26.
50. Harvey RL, Edwards D, Dunning K, et al. Randomized Sham-Controlled Trial of Navigated Repetitive Transcranial Magnetic Stimulation for Motor Recovery in Stroke. *Stroke.* 2018;49:2138-2146.
51. Edwards DJ, Liu CY, Dunning K, et al. Electric Field Navigated 1-Hz rTMS for Poststroke Motor Recovery: The E-FIT Randomized Controlled Trial. *Stroke.* 2023;54:2254-2264.
52. Chen G, Lin T, Wu M, et al. Effects of repetitive transcranial magnetic stimulation on upper-limb and finger function in stroke patients: A systematic review and meta-analysis of randomized controlled trials. *Front Neurol.* 2022;13:940467.
53. Mead G, Graham C, Lundström E, et al. Individual patient data meta-analysis of the effects of fluoxetine on functional outcomes after acute stroke. *Int J Stroke.* 2024;19:798-808.
54. Safety and efficacy of fluoxetine on functional recovery after acute stroke (EFFECTS): a randomised, double-blind, placebo-controlled trial. *The Lancet. Neurology.* 2020;19:661-669.
55. Safety and efficacy of fluoxetine on functional outcome after acute stroke (AFFINITY): a randomised, double-blind, placebo-controlled trial. *The Lancet. Neurology.* 2020;19:651-660.

56. Effects of fluoxetine on functional outcomes after acute stroke (FOCUS): a pragmatic, double-blind, randomised, controlled trial. *Lancet*. 2019;393:265-274.
57. Legg LA, Rudberg AS, Hua X, et al. Selective serotonin reuptake inhibitors (SSRIs) for stroke recovery. *Cochrane Database Syst Rev*. 2021;11:Cd009286.
58. Petrea RE, Beiser AS, Seshadri S, Kelly-Hayes M, Kase CS, Wolf PA. Stroke in women-Gender differences in stroke incidence and post-stroke disability in the Framingham Heart Study. *Stroke*. 2009;40:1032–1037.
59. MacDonald SL, Hall RE, Bell CM, Cronin S, Jaglal SB. Sex differences in the outcomes of adults admitted to inpatient rehabilitation after stroke. *PM R*. 2022;14:779-785.
60. Mehrabi S, Harnett A, Saikaley M, et al. Female Enrollment in Rehabilitation Trials: A Systematic Review of Reporting Sex and Female Participation in Randomized Controlled Trials of Poststroke Upper Extremity Rehabilitation Over 50 Years. *Arch Phys Med Rehabil*. 2024;105:1399-1406.
61. Anwer S, Alghadir A. Incidence, Prevalence, and Risk Factors of Hemiplegic Shoulder Pain: A Systematic Review. *Int J Environ Res Public Health*. 2020;17.
62. Ada L, Foongchomcheay A, Langhammer B, et al. Lap-tray and triangular sling are no more effective than a hemi-sling in preventing shoulder subluxation in those at risk early after stroke: a randomized trial. *Eur J Phys Rehabil Med*. 2017;53:41-48.
63. Pan R, Zhou M, Cai H, et al. A randomized controlled trial of a modified wheelchair arm-support to reduce shoulder pain in stroke patients. *Clin Rehabil*. 2018;32:37-47.
64. Deng P, Zhao Z, Zhang S, Xiao T, Li Y. Effect of kinesio taping on hemiplegic shoulder pain: A systematic review and meta-analysis of randomized controlled trials. *Clin Rehabil*. 2021;35:317-331.
65. Appel C, Perry L, Jones F. Shoulder strapping for stroke-related upper limb dysfunction and shoulder impairments: systematic review. *NeuroRehabilitation*. 2014;35:191-204.
66. Ada L, Foongchomcheay A, Canning C. Supportive devices for preventing and treating subluxation of the shoulder after stroke. *Cochrane Database Syst Rev*. 2005;2005:Cd003863.
67. Qiu H, Li J, Zhou T, Wang H, Li J. Electrical Stimulation in the Treatment of Hemiplegic Shoulder Pain: A Meta-Analysis of Randomized Controlled Trials. *Am J Phys Med Rehabil*. 2019;98:280-286.
68. Lee JH, Baker LL, Johnson RE, Tilson JK. Effectiveness of neuromuscular electrical stimulation for management of shoulder subluxation post-stroke: a systematic review with meta-analysis. *Clin Rehabil*. 2017;31:1431-1444.
69. Church C, Price C, Pandyan AD, Huntley S, Curless R, Rodgers H. Randomized controlled trial to evaluate the effect of surface neuromuscular electrical stimulation to the shoulder after acute stroke. *Stroke*. 2006;37:2995-3001.

70. Singh JA, Fitzgerald PM. Botulinum toxin for shoulder pain. *Cochrane Database Syst Rev*. 2010;Cd008271.
71. de Boer KS, Arwert HJ, de Groot JH, Meskers CG, Mishre AD, Arendzen JH. Shoulder pain and external rotation in spastic hemiplegia do not improve by injection of botulinum toxin A into the subscapular muscle. *J Neurol Neurosurg Psychiatry*. 2008;79:581-583.
72. Tan B, Jia L. Ultrasound-Guided BoNT-A (Botulinum Toxin A) Injection Into the Subscapularis for Hemiplegic Shoulder Pain: A Randomized, Double-Blind, Placebo-Controlled Trial. *Stroke*. 2021;52:3759-3767.
73. Rah UW, Yoon SH, Moon DJ, et al. Subacromial corticosteroid injection on poststroke hemiplegic shoulder pain: a randomized, triple-blind, placebo-controlled trial. *Arch Phys Med Rehabil*. 2012;93:949-956.
74. Snels IA, Beckerman H, Twisk JW, et al. Effect of triamcinolone acetonide injections on hemiplegic shoulder pain : A randomized clinical trial. *Stroke*. 2000;31:2396-2401.
75. Zhan J, Wei X, Tao C, et al. Effectiveness of acupuncture combined with rehabilitation training vs. rehabilitation training alone for post-stroke shoulder pain: A systematic review and meta-analysis of randomized controlled trials. *Front Med (Lausanne)*. 2022;9:947285.
76. Zhang T, Zhang C. Extracorporeal shock wave therapy for shoulder pain after stroke: A systematic review and meta-analysis. *Clin Rehabil*. 2023;37:774-790.
77. Bhat C, Rosenberg H, James D. Topical nonsteroidal anti-inflammatory drugs. *CMAJ*. 2023;195:E1231.
78. Giang TA, Ong AWG, Krishnamurthy K, Fong KNK. Rehabilitation Interventions for Poststroke Hand Oedema: A Systematic Review. *Hong Kong J Occup Ther*. 2016;27:7-17.
79. O'Connell NE, Wand BM, McAuley J, Marston L, Moseley GL. Interventions for treating pain and disability in adults with complex regional pain syndrome. *Cochrane Database Syst Rev*. 2013;2013:Cd009416.
80. Liu S, Zhang CS, Cai Y, et al. Acupuncture for Post-stroke Shoulder-Hand Syndrome: A Systematic Review and Meta-Analysis. *Front Neurol*. 2019;10:433.
81. Peng L, Zhang C, Zhou L, Zuo HX, He XK, Niu YM. Traditional manual acupuncture combined with rehabilitation therapy for shoulder hand syndrome after stroke within the Chinese healthcare system: a systematic review and meta-analysis. *Clin Rehabil*. 2018;32:429-439.
82. Harvey LA, Katalinic OM, Herbert RD, Moseley AM, Lannin NA, Schurr K. Stretch for the treatment and prevention of contracture: an abridged republication of a Cochrane Systematic Review. *J Physiother*. 2017;63:67-75.



83. Salazar AP, Pinto C, Ruschel Mossi JV, Figueiro B, Lukrafka JL, Pagnussat AS. Effectiveness of static stretching positioning on post-stroke upper-limb spasticity and mobility: Systematic review with meta-analysis. *Ann Phys Rehabil Med*. 2019;62:274-282.
84. Ward AB, Wissel J, Borg J, et al. Functional goal achievement in post-stroke spasticity patients: the BOTOX(R) Economic Spasticity Trial (BEST). *J Rehabil Med*. 2014;46:504-513.
85. Wissel J, Ganapathy V, Ward AB, et al. OnabotulinumtoxinA Improves Pain in Patients With Post-Stroke Spasticity: Findings From a Randomized, Double-Blind, Placebo-Controlled Trial. *J Pain Symptom Manage*. 2016;52:17-26.
86. Shaw LC, Price CI, van Wijck FM, et al. Botulinum Toxin for the Upper Limb after Stroke (BoTULS) Trial: effect on impairment, activity limitation, and pain. *Stroke*. 2011;42:1371-1379.
87. McCrory P, Turner-Stokes L, Baguley IJ, et al. Botulinum toxin A for treatment of upper limb spasticity following stroke: a multi-centre randomized placebo-controlled study of the effects on quality of life and other person-centred outcomes. *J Rehabil Med*. 2009;41:536-544.
88. Sun LC, Chen R, Fu C, et al. Efficacy and Safety of Botulinum Toxin Type A for Limb Spasticity after Stroke: A Meta-Analysis of Randomized Controlled Trials. *Biomed Res Int*. 2019;2019:8329306.
89. Lee JM, Gracies JM, Park SB, Lee KH, Lee JY, Shin JH. Botulinum Toxin Injections and Electrical Stimulation for Spastic Paresis Improve Active Hand Function Following Stroke. *Toxins (Basel)*. 2018;10.
90. Santamato A, Notarnicola A, Panza F, et al. SBOTE study: extracorporeal shock wave therapy versus electrical stimulation after botulinum toxin type a injection for post-stroke spasticity-a prospective randomized trial. *Ultrasound Med Biol*. 2013;39:283-291.
91. Nasb M, Li Z, A SAY, Dayoub L, Chen H. Comparison of the effects of modified constraint-induced movement therapy and intensive conventional therapy with a botulinum-a toxin injection on upper limb motor function recovery in patients with stroke. *Libyan J Med*. 2019;14:1609304.
92. Santamato A, Micello MF, Panza F, et al. Adhesive taping vs. daily manual muscle stretching and splinting after botulinum toxin type A injection for wrist and fingers spastic overactivity in stroke patients: a randomized controlled trial. *Clin Rehabil*. 2015;29:50-58.
93. Amini M, Shamili A, Frough B, Pashmdarfard M, Fallahzadeh Abarghouei A. Combined effect of botulinum toxin and splinting on motor components and function of people suffering a stroke. *Med J Islam Repub Iran*. 2016;30:373.
94. Güntürk E, Ögüt H, Güler H, Turhanoğlu AD. The effect of oral baclofen and botulinum toxin treatments in hemiplegic spasticity on the nociceptive flexor reflex: A randomized clinical trial. *Turk J Phys Med Rehabil*. 2022;68:524-531.
95. Gelber DA, Good DC, Dromerick A, Sergay S, Richardson M. Open-label dose-titration safety and efficacy study of tizanidine hydrochloride in the treatment of spasticity associated with chronic stroke. *Stroke*. 2001;32:1841-1846.



96. Cabanas-Valdés R, Serra-Llobet P, Rodriguez-Rubio PR, López-de-Celis C, Llauro-Fores M, Calvo-Sanz J. The effectiveness of extracorporeal shock wave therapy for improving upper limb spasticity and functionality in stroke patients: a systematic review and meta-analysis. *Clin Rehabil*. 2020;34:1141-1156.
97. Wang X, Ge L, Hu H, Yan L, Li L. Effects of Non-Invasive Brain Stimulation on Post-Stroke Spasticity: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Brain Sci*. 2022;12.
98. Chen S, Lv C, Wu J, Zhou C, Shui X, Wang Y. Effectiveness of a home-based exercise program among patients with lower limb spasticity post-stroke: A randomized controlled trial. *Asian Nurs Res (Korean Soc Nurs Sci)*. 2021;15:1-7.
99. Kluding PM, Santos M. Effects of ankle joint mobilizations in adults poststroke: a pilot study. *Arch Phys Med Rehabil*. 2008;89:449-456.
100. Wein T, Esquenazi A, Jost WH, Ward AB, Pan G, Dimitrova R. OnabotulinumtoxinA for the Treatment of Poststroke Distal Lower Limb Spasticity: A Randomized Trial. *PM R*. 2018;10:693-703.
101. Doan TN, Kuo MY, Chou LW. Efficacy and Optimal Dose of Botulinum Toxin A in Post-Stroke Lower Extremity Spasticity: A Systematic Review and Meta-Analysis. *Toxins (Basel)*. 2021;13.
102. Kaji R, Osako Y, Suyama K, Maeda T, Uechi Y, Iwasaki M. Botulinum toxin type A in post-stroke lower limb spasticity: a multicenter, double-blind, placebo-controlled trial. *J Neurol*. 2010;257:1330-1337.
103. Pittock SJ, Moore AP, Hardiman O, et al. A double-blind randomised placebo-controlled evaluation of three doses of botulinum toxin type A (Dysport) in the treatment of spastic equinovarus deformity after stroke. *Cerebrovasc Dis*. 2003;15:289-300.
104. Creamer M, Cloud G, Kossmehl P, et al. Effect of Intrathecal Baclofen on Pain and Quality of Life in Poststroke Spasticity. *Stroke*. 2018;49:2129-2137.
105. Yoldaş Aslan Ş, Kutlay S, Düsünceli Atman E, Elhan AH, Gök H, Küçükdeveci AA. Does extracorporeal shock wave therapy decrease spasticity of ankle plantar flexor muscles in patients with stroke: A randomized controlled trial. *Clin Rehabil*. 2021;35:1442-1453.
106. Lee CH, Lee SH, Yoo JI, Lee SU. Ultrasonographic Evaluation for the Effect of Extracorporeal Shock Wave Therapy on Gastrocnemius Muscle Spasticity in Patients With Chronic Stroke. *PM R*. 2019;11:363-371.
107. MacKay-Lyons M, Billinger SA, Eng JJ, et al. Aerobic Exercise Recommendations to Optimize Best Practices in Care After Stroke: AEROBICS 2019 Update. *Phys Ther*. 2020;100:149-156.
108. Van Crielinge T, Heremans C, Burridge J, et al. Standardized measurement of balance and mobility post-stroke: Consensus-based core recommendations from the third Stroke Recovery and Rehabilitation Roundtable. *Int J Stroke*. 2024;19:158-168.

109. English C, Hillier SL, Lynch EA. Circuit class therapy for improving mobility after stroke. *Cochrane Database Syst Rev*. 2017;6:Cd007513.
110. Flansbjerg UB, Miller M, Downham D, Lexell J. Progressive resistance training after stroke: effects on muscle strength, muscle tone, gait performance and perceived participation. *J Rehabil Med*. 2008;40:42-48.
111. Flansbjerg UB, Lexell J, Brogårdh C. Long-term benefits of progressive resistance training in chronic stroke: a 4-year follow-up. *J Rehabil Med*. 2012;44:218-221.
112. Cooke EV, Tallis RC, Clark A, Pomeroy VM. Efficacy of functional strength training on restoration of lower-limb motor function early after stroke: phase I randomized controlled trial. *Neurorehabil Neural Repair*. 2010;24:88-96.
113. Mehrholz J, Thomas S, Elsner B. Treadmill training and body weight support for walking after stroke. *Cochrane Database Syst Rev*. 2017;8:Cd002840.
114. Ada L, Dean CM, Morris ME, Simpson JM, Katrak P. Randomized trial of treadmill walking with body weight support to establish walking in subacute stroke: the MOBILISE trial. *Stroke*. 2010;41:1237-1242.
115. Dean CM, Ada L, Bampton J, Morris ME, Katrak PH, Potts S. Treadmill walking with body weight support in subacute non-ambulatory stroke improves walking capacity more than overground walking: a randomised trial. *J Physiother*. 2010;56:97-103.
116. Nadeau SE, Wu SS, Dobkin BH, et al. Effects of task-specific and impairment-based training compared with usual care on functional walking ability after inpatient stroke rehabilitation: LEAPS Trial. *Neurorehabil Neural Repair*. 2013;27:370-380.
117. Mehrholz J, Thomas S, Kugler J, Pohl M, Elsner B. Electromechanical-assisted training for walking after stroke. *Cochrane Database Syst Rev*. 2020;10:Cd006185.
118. Molteni F, Guanzirolì E, Goffredo M, et al. Gait Recovery with an Overground Powered Exoskeleton: A Randomized Controlled Trial on Subacute Stroke Subjects. *Brain Sci*. 2021;11.
119. Ghai S, Ghai I. Effects of (music-based) rhythmic auditory cueing training on gait and posture post-stroke: A systematic review & dose-response meta-analysis. *Sci Rep*. 2019;9:2183.
120. Yoo GE, Kim SJ. Rhythmic Auditory Cueing in Motor Rehabilitation for Stroke Patients: Systematic Review and Meta-Analysis. *J Music Ther*. 2016;53:149-177.
121. Zhang B, Li D, Liu Y, Wang J, Xiao Q. Virtual reality for limb motor function, balance, gait, cognition and daily function of stroke patients: A systematic review and meta-analysis. *J Adv Nurs*. 2021;77:3255-3273.
122. Iruthayarajah J, McIntyre A, Cotoi A, Macaluso S, Teasell R. The use of virtual reality for balance among individuals with chronic stroke: a systematic review and meta-analysis. *Top Stroke Rehabil*. 2017;24:68-79.

123. Gibbons EM, Thomson AN, de Noronha M, Joseph S. Are virtual reality technologies effective in improving lower limb outcomes for patients following stroke - a systematic review with meta-analysis. *Top Stroke Rehabil.* 2016;23:440-457.
124. Silva S, Borges LR, Santiago L, Lucena L, Lindquist AR, Ribeiro T. Motor imagery for gait rehabilitation after stroke. *Cochrane Database Syst Rev.* 2020;9:Cd013019.
125. Jaqueline da Cunha M, Rech KD, Salazar AP, Pagnussat AS. Functional electrical stimulation of the peroneal nerve improves post-stroke gait speed when combined with physiotherapy. A systematic review and meta-analysis. *Ann Phys Rehabil Med.* 2021;64:101388.
126. Howlett OA, Lannin NA, Ada L, McKinstry C. Functional electrical stimulation improves activity after stroke: a systematic review with meta-analysis. *Arch Phys Med Rehabil.* 2015;96:934-943.
127. Pomeroy VM, King L, Pollock A, Baily-Hallam A, Langhorne P. Electrostimulation for promoting recovery of movement or functional ability after stroke. *Cochrane Database Syst Rev.* 2006;2006:Cd003241.
128. Stanton R, Ada L, Dean CM, Preston E. Biofeedback improves performance in lower limb activities more than usual therapy in people following stroke: a systematic review. *J Physiother.* 2017;63:11-16.
129. Daryabor A, Kobayashi T, Yamamoto S, Lyons SM, Orendurff M, Akbarzadeh Baghban A. Effect of ankle-foot orthoses on functional outcome measurements in individuals with stroke: a systematic review and meta-analysis. *Disabil Rehabil.* 2022;44:6566-6581.
130. Choo YJ, Chang MC. Effectiveness of an ankle-foot orthosis on walking in patients with stroke: a systematic review and meta-analysis. *Sci Rep.* 2021;11:15879.
131. Tyson SF, Kent RM. Effects of an ankle-foot orthosis on balance and walking after stroke: a systematic review and pooled meta-analysis. *Arch Phys Med Rehabil.* 2013;94:1377-1385.
132. Erel S, Uygur F, Engin Simsek I, Yakut Y. The effects of dynamic ankle-foot orthoses in chronic stroke patients at three-month follow-up: a randomized controlled trial. *Clin Rehabil.* 2011;25:515-523.
133. Ghayour Najafabadi M, Shariat A, Dommerholt J, et al. Aquatic Therapy for improving Lower Limbs Function in Post-stroke Survivors: A Systematic Review with Meta-Analysis. *Top Stroke Rehabil.* 2022;29:473-489.
134. Chae CS, Jun JH, Im S, Jang Y, Park GY. Effectiveness of Hydrotherapy on Balance and Paretic Knee Strength in Patients With Stroke: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Am J Phys Med Rehabil.* 2020;99:409-419.
135. Bank J, Charles K, Morgan P. What is the effect of additional physiotherapy on sitting balance following stroke compared to standard physiotherapy treatment: a systematic review. *Top Stroke Rehabil.* 2016;23:15-25.

136. Barclay-Goddard R, Stevenson T, Poluha W, Moffatt ME, Taback SP. Force platform feedback for standing balance training after stroke. *Cochrane Database Syst Rev*. 2004;2004:Cd004129.
137. Van Peppen RP, Kortsmit M, Lindeman E, Kwakkel G. Effects of visual feedback therapy on postural control in bilateral standing after stroke: a systematic review. *J Rehabil Med*. 2006;38:3-9.
138. Saquette MB, da Silva CM, Martinez BP, et al. Water-Based Exercise on Functioning and Quality of Life in Poststroke Persons: A Systematic Review and Meta-Analysis. *J Stroke Cerebrovasc Dis*. 2019;28:104341.
139. Zhang L, Zhang L, Yu X, Zhou H, Ding Y, Wang J. Effect of Tai Chi Yunshou training on the balance and motor functions of stroke patients: a systematic review and meta-analysis of randomized controlled trials. *Front Neurol*. 2023;14:1178234.
140. Zheng X, Wu X, Liu Z, et al. The Influences of Tai Chi on Balance Function and Exercise Capacity among Stroke Patients: A Meta-Analysis. *Evid Based Complement Alternat Med*. 2021;2021:6636847.
141. Ge L, Zheng QX, Liao YT, Tan JY, Xie QL, Rask M. Effects of traditional Chinese exercises on the rehabilitation of limb function among stroke patients: A systematic review and meta-analysis. *Complement Ther Clin Pract*. 2017;29:35-47.
142. Zhao LJ, Jiang LH, Zhang H, et al. Effects of Motor Imagery Training for Lower Limb Dysfunction in Patients With Stroke: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Am J Phys Med Rehabil*. 2023;102:409-418.
143. Yin Y, Wang J, Yu Z, et al. Does whole-body vibration training have a positive effect on balance and walking function in patients with stroke? A meta-analysis. *Front Hum Neurosci*. 2022;16:1076665.
144. Yang X, Xue X, Tu H, Li N. Effect of whole-body vibration training on the recovery of lower limb function in people with stroke: a systematic review and meta-analysis. *Disabil Rehabil*. 2023;45:3823-3832.
145. Pollock A, Gray C, Culham E, Durward BR, Langhorne P. Interventions for improving sit-to-stand ability following stroke. *Cochrane Database Syst Rev*. 2014;2014:Cd007232.
146. Saunders DH, Sanderson M, Hayes S, et al. Physical fitness training for stroke patients. *Cochrane Database Syst Rev*. 2020;3:Cd003316.
147. Sandberg K, Kleist M, Falk L, Enthoven P. Effects of Twice-Weekly Intense Aerobic Exercise in Early Subacute Stroke: A Randomized Controlled Trial. *Arch Phys Med Rehabil*. 2016;97:1244-1253.
148. Hornby TG, Holleran CL, Hennessy PW, et al. Variable Intensive Early Walking Poststroke (VIEWS): A Randomized Controlled Trial. *Neurorehabil Neural Repair*. 2016;30:440-450.

149. Jin H, Jiang Y, Wei Q, Wang B, Ma G. Intensive aerobic cycling training with lower limb weights in Chinese patients with chronic stroke: discordance between improved cardiovascular fitness and walking ability. *Disabil Rehabil.* 2012;34:1665-1671.
150. Globas C, Becker C, Cerny J, et al. Chronic stroke survivors benefit from high-intensity aerobic treadmill exercise: a randomized control trial. *Neurorehabil Neural Repair.* 2012;26:85-95.
151. Safaei-Qomi MR, Mehrabi S, Fleet JL, et al. A Systematic Review of Worldwide Female Enrollment in Randomized Controlled Trials of Post-Stroke Lower Extremity Rehabilitation. *Am J Phys Med Rehabil.* 2024.
152. Zhang L, Wang J, Dong X, et al. Injurious Falls before, during, and after Stroke Diagnosis: A Population-based Study. *Journal of the American Medical Directors Association.* 2025;26:105465.
153. Czernuszenko A, Czlonkowska A. Risk factors for falls in stroke patients during inpatient rehabilitation. *Clin Rehabil.* 2009;23:176-188.
154. Xu T, Clemson L, O'Loughlin K, Lannin NA, Dean C, Koh G. Risk Factors for Falls in Community Stroke Survivors: A Systematic Review and Meta-Analysis. *Arch Phys Med Rehabil.* 2018;99:563-573.e565.
155. Strini V, Schiavolin R, Prendin A. Fall Risk Assessment Scales: A Systematic Literature Review. *Nurs Rep.* 2021;11:430-443.
156. Breisinger TP, Skidmore ER, Niyonkuru C, Terhorst L, Campbell GB. The Stroke Assessment of Fall Risk (SAFR): predictive validity in inpatient stroke rehabilitation. *Clin Rehabil.* 2014;28:1218-1224.
157. Nyström A, Hellström K. Fall risk six weeks from onset of stroke and the ability of the Prediction of Falls in Rehabilitation Settings Tool and motor function to predict falls. *Clin Rehabil.* 2013;27:473-479.
158. Pinto EB, Nascimento C, Monteiro M, et al. Proposal for a New Predictive Scale for Recurrent Risk of Fall in a Cohort of Community-Dwelling Patients with Stroke. *J Stroke Cerebrovasc Dis.* 2016;25:2619-2626.
159. Mansfield A, Aqui A, Danells CJ, et al. Does perturbation-based balance training prevent falls among individuals with chronic stroke? A randomised controlled trial. *BMJ Open.* 2018;8:e021510.
160. Dean CM, Rissel C, Sherrington C, et al. Exercise to enhance mobility and prevent falls after stroke: the community stroke club randomized trial. *Neurorehabil Neural Repair.* 2012;26:1046-1057.
161. Batchelor FA, Hill KD, Mackintosh SF, Said CM, Whitehead CH. Effects of a multifactorial falls prevention program for people with stroke returning home after rehabilitation: a randomized controlled trial. *Arch Phys Med Rehabil.* 2012;93:1648-1655.
162. Denissen S, Staring W, Kunkel D, et al. Interventions for preventing falls in people after stroke. *Cochrane Database Syst Rev.* 2019;10:Cd008728.

163. Yang F, Lees J, Simpkins C, Butler A. Interventions for preventing falls in people post-stroke: A meta-analysis of randomized controlled trials. *Gait Posture*. 2021;84:377-388.
164. Martino R, Foley N, Bhogal S, Diamant N, Speechley M, Teasell R. Dysphagia after stroke: incidence, diagnosis, and pulmonary complications. *Stroke*. 2005;36:2756-2763.
165. Boaden E, Burnell J, Hives L, et al. Screening for aspiration risk associated with dysphagia in acute stroke. *Cochrane Database Syst Rev*. 2021;10:Cd012679.
166. Trapl M, Enderle P, Nowotny M, et al. Dysphagia bedside screening for acute-stroke patients: the Gugging Swallowing Screen. *Stroke*. 2007;38:2948-2952.
167. Martino R, Silver F, Teasell R, et al. The Toronto Bedside Swallowing Screening Test (TOR-BSST): development and validation of a dysphagia screening tool for patients with stroke. *Stroke*. 2009;40:555-561.
168. Sherman V, Greco E, Martino R. The Benefit of Dysphagia Screening in Adult Patients With Stroke: A Meta-Analysis. *J Am Heart Assoc*. 2021;10:e018753.
169. Middleton S, McElduff P, Ward J, et al. Implementation of evidence-based treatment protocols to manage fever, hyperglycaemia, and swallowing dysfunction in acute stroke (QASC): a cluster randomised controlled trial. *Lancet*. 2011;378:1699-1706.
170. Tohara H, Saitoh E, Mays KA, Kuhlemeier K, Palmer JB. Three tests for predicting aspiration without videofluorography. *Dysphagia*. 2003;18:126-134.
171. Wu MC, Chang YC, Wang TG, Lin LC. Evaluating swallowing dysfunction using a 100-ml water swallowing test. *Dysphagia*. 2004;19:43-47.
172. Schrock JW, Bernstein J, Glasenapp M, Drogell K, Hanna J. A novel emergency department dysphagia screen for patients presenting with acute stroke. *Acad Emerg Med*. 2011;18:584-589.
173. Bath PM, Lee HS, Everton LF. Swallowing therapy for dysphagia in acute and subacute stroke. *Cochrane Database Syst Rev*. 2018;10:Cd000323.
174. Duncan S, McAuley DF, Walshe M, et al. Interventions for oropharyngeal dysphagia in acute and critical care: a systematic review and meta-analysis. *Intensive Care Med*. 2020;46:1326-1338.
175. Tarihci Cakmak E, Sen EI, Doruk C, Sen C, Sezikli S, Yaliman A. The Effects of Neuromuscular Electrical Stimulation on Swallowing Functions in Post-stroke Dysphagia: A Randomized Controlled Trial. *Dysphagia*. 2023;38:874-885.
176. Carnaby GD, LaGorio L, Silliman S, Crary M. Exercise-based swallowing intervention (McNeill Dysphagia Therapy) with adjunctive NMES to treat dysphagia post-stroke: A double-blind placebo-controlled trial. *J Oral Rehabil*. 2020;47:501-510.



177. Park JS, Oh DH, Hwang NK, Lee JH. Effects of neuromuscular electrical stimulation combined with effortful swallowing on post-stroke oropharyngeal dysphagia: a randomised controlled trial. *J Oral Rehabil.* 2016;43:426-434.
178. Chen YW, Chang KH, Chen HC, Liang WM, Wang YH, Lin YN. The effects of surface neuromuscular electrical stimulation on post-stroke dysphagia: a systemic review and meta-analysis. *Clin Rehabil.* 2016;30:24-35.
179. Dennis MS, Lewis SC, Warlow C. Effect of timing and method of enteral tube feeding for dysphagic stroke patients (FOOD): a multicentre randomised controlled trial. *Lancet.* 2005;365:764-772.
180. Gomes CA, Jr., Andriolo RB, Bennett C, et al. Percutaneous endoscopic gastrostomy versus nasogastric tube feeding for adults with swallowing disturbances. *Cochrane Database Syst Rev.* 2015:Cd008096.
181. Deutz NE, Matheson EM, Matarese LE, et al. Readmission and mortality in malnourished, older, hospitalized adults treated with a specialized oral nutritional supplement: A randomized clinical trial. *Clin Nutr.* 2016;35:18-26.
182. Schuetz P, Fehr R, Baechli V, et al. Individualised nutritional support in medical inpatients at nutritional risk: a randomised clinical trial. *Lancet.* 2019;393:2312-2321.
183. Dennis MS, Lewis SC, Warlow C. Routine oral nutritional supplementation for stroke patients in hospital (FOOD): a multicentre randomised controlled trial. *Lancet.* 2005;365:755-763.
184. Sakai K, Niimi M, Momosaki R, et al. Nutritional therapy for reducing disability and improving activities of daily living in people after stroke. *Cochrane Database Syst Rev.* 2024;8:Cd014852.
185. Campbell P, Bain B, Furlanetto DL, Brady MC. Interventions for improving oral health in people after stroke. *Cochrane Database Syst Rev.* 2020;12:Cd003864.
186. Pedersen PM, Jørgensen HS, Nakayama H, Raaschou HO, Olsen TS. Aphasia in acute stroke: incidence, determinants, and recovery. *Ann Neurol.* 1995;38:659-666.
187. Brust JC, Shafer SQ, Richter RW, Bruun B. Aphasia in acute stroke. *Stroke.* 1976;7:167-174.
188. Lazar RM, Boehme AK. Aphasia As a Predictor of Stroke Outcome. *Curr Neurol Neurosci Rep.* 2017;17:83.
189. Brady MC, Kelly H, Godwin J, Enderby P, Campbell P. Speech and language therapy for aphasia following stroke. *Cochrane Database Syst Rev.* 2016:Cd000425.
190. Dosage, Intensity, and Frequency of Language Therapy for Aphasia: A Systematic Review-Based, Individual Participant Data Network Meta-Analysis. *Stroke.* 2022;53:956-967.
191. Simmons-Mackie N, Raymer A, Cherney LR. Communication Partner Training in Aphasia: An Updated Systematic Review. *Arch Phys Med Rehabil.* 2016;97:2202-2221.e2208.



192. Simmons-Mackie N, Raymer A, Armstrong E, Holland A, Cherney LR. Communication partner training in aphasia: a systematic review. *Arch Phys Med Rehabil*. 2010;91:1814-1837.
193. Li TT, Zhang PP, Zhang MC, et al. Meta-analysis and systematic review of the relationship between sex and the risk or incidence of poststroke aphasia and its types. *BMC Geriatr*. 2024;24:220.
194. Wallentin M. Sex differences in post-stroke aphasia rates are caused by age. A meta-analysis and database query. *PLoS One*. 2018;13:e0209571.
195. Brady MC, Ali M, VandenBerg K, et al. Precision rehabilitation for aphasia by patient age, sex, aphasia severity, and time since stroke? A prespecified, systematic review-based, individual participant data, network, subgroup meta-analysis. *Int J Stroke*. 2022;17:1067-1077.
196. Rowe FJ, Hepworth LR, Howard C, Hanna KL, Cheyne CP, Currie J. High incidence and prevalence of visual problems after acute stroke: An epidemiology study with implications for service delivery. *PLoS One*. 2019;14:e0213035.
197. Rowe FJ. Stroke survivors' views and experiences on impact of visual impairment. *Brain Behav*. 2017;7:e00778.
198. Hepworth LR, Howard C, Hanna KL, Currie J, Rowe FJ. "Eye" Don't See: An Analysis of Visual Symptom Reporting by Stroke Survivors from a Large Epidemiology Study. *J Stroke Cerebrovasc Dis*. 2021;30:105759.
199. Rowe FJ, Hepworth LR, Kirkham JJ. Development of core outcome sets for vision screening and assessment in stroke: a Delphi and consensus study. *BMJ Open*. 2019;9:e029578.
200. Donkervoort M, Dekker J, van den Ende E, Stehmann-Saris JC, Deelman BG. Prevalence of apraxia among patients with a first left hemisphere stroke in rehabilitation centres and nursing homes. *Clin Rehabil*. 2000;14:130-136.
201. Hepworth L, Rowe F, Walker M, et al. Post-stroke visual impairment: a systematic literature review of types and recovery of visual conditions. *Ophthalmol Res*. 2016;5:1-43.
202. Hazelton C, Thomson K, Todhunter-Brown A, et al. Interventions for perceptual disorders following stroke. *Cochrane Database Syst Rev*. 2022;11:Cd007039.
203. Longley V, Hazelton C, Heal C, et al. Non-pharmacological interventions for spatial neglect or inattention following stroke and other non-progressive brain injury. *Cochrane Database Syst Rev*. 2021;7:Cd003586.
204. Pollock A, Hazelton C, Rowe FJ, et al. Interventions for visual field defects in people with stroke. *Cochrane Database Syst Rev*. 2019;5:Cd008388.
205. Zhang Y, Xing Y, Li C, et al. Mirror therapy for unilateral neglect after stroke: A systematic review. *Eur J Neurol*. 2022;29:358-371.

206. Aparicio-López C, García-Molina A, García-Fernández J, et al. Cognitive rehabilitation with right hemifield eye-patching for patients with sub-acute stroke and visuo-spatial neglect: a randomized controlled trial. *Brain Inj.* 2015;29:501-507.
207. Tsang MH, Sze KH, Fong KN. Occupational therapy treatment with right half-field eye-patching for patients with subacute stroke and unilateral neglect: a randomised controlled trial. *Disabil Rehabil.* 2009;31:630-637.
208. Kim YM, Chun MH, Yun GJ, Song YJ, Young HE. The effect of virtual reality training on unilateral spatial neglect in stroke patients. *Ann Rehabil Med.* 2011;35:309-315.
209. Katz N, Ring H, Naveh Y, Kizony R, Feintuch U, Weiss PL. Interactive virtual environment training for safe street crossing of right hemisphere stroke patients with unilateral spatial neglect. *Disabil Rehabil.* 2005;27:1235-1243.
210. Luukkainen-Markkula R, Tarkka IM, Pitkänen K, Sivenius J, Hämäläinen H. Rehabilitation of hemispatial neglect: A randomized study using either arm activation or visual scanning training. *Restor Neurol Neurosci.* 2009;27:663-672.
211. Kim BR, Chun MH, Kim DY, Lee SJ. Effect of high- and low-frequency repetitive transcranial magnetic stimulation on visuospatial neglect in patients with acute stroke: a double-blind, sham-controlled trial. *Arch Phys Med Rehabil.* 2013;94:803-807.
212. Yang NY, Fong KN, Li-Tsang CW, Zhou D. Effects of repetitive transcranial magnetic stimulation combined with sensory cueing on unilateral neglect in subacute patients with right hemispheric stroke: a randomized controlled study. *Clin Rehabil.* 2017;31:1154-1163.
213. Andersen G, Vestergaard K, Ingeman-Nielsen M, Jensen TS. Incidence of central post-stroke pain. *Pain.* 1995;61:187-193.
214. Mahesh B, Singh VK, Pathak A, et al. Efficacy of Duloxetine in Patients with Central Post-stroke Pain: A Randomized Double Blind Placebo Controlled Trial. *Pain Med.* 2023;24:610-617.
215. Vranken JH, Hollmann MW, van der Vegt MH, et al. Duloxetine in patients with central neuropathic pain caused by spinal cord injury or stroke: a randomized, double-blind, placebo-controlled trial. *Pain.* 2011;152:267-273.
216. Kim NY, Lee SC, Kim YW. Effect of Duloxetine for the Treatment of Chronic Central Poststroke Pain. *Clin Neuropharmacol.* 2019;42:73-76.
217. Kim JS, Bashford G, Murphy KT, Martin A, Dror V, Cheung R. Safety and efficacy of pregabalin in patients with central post-stroke pain. *Pain.* 2011;152:1018-1023.
218. Vranken JH, Dijkgraaf MG, Kruis MR, van der Vegt MH, Hollmann MW, Heesen M. Pregabalin in patients with central neuropathic pain: a randomized, double-blind, placebo-controlled trial of a flexible-dose regimen. *Pain.* 2008;136:150-157.

219. Serpell MG. Gabapentin in neuropathic pain syndromes: a randomised, double-blind, placebo-controlled trial. *Pain*. 2002;99:557-566.
220. Jungehulsing GJ, Israel H, Safar N, et al. Levetiracetam in patients with central neuropathic post-stroke pain--a randomized, double-blind, placebo-controlled trial. *Eur J Neurol*. 2013;20:331-337.
221. Thomas LH, Coupe J, Cross LD, Tan AL, Watkins CL. Interventions for treating urinary incontinence after stroke in adults. *Cochrane Database Syst Rev*. 2019;2:Cd004462.
222. Özden F, Tümtürk İ, Özkeskin M, Bakırhan S. The effect of pelvic floor muscle training on urinary incontinence in patients with stroke: a systematic review and meta-analysis. *Ir J Med Sci*. 2023;192:1481-1495.
223. Cruz E, Miller C, Zhang W, et al. Does non-implanted electrical stimulation reduce post-stroke urinary or fecal incontinence? A systematic review with meta-analysis. *Int J Stroke*. 2022;17:378-388.
224. Thomas LH, Watkins CL, Sutton CJ, et al. Identifying continence options after stroke (ICONS): a cluster randomised controlled feasibility trial. *Trials*. 2014;15:509.
225. Watkins C, Tishkovskaya S, Brown C, et al. Systematic voiding programme in adults with urinary incontinence following acute stroke: the ICONS-II RCT. *Health Technol Assess*. 2022;26:1-88.
226. Tibaek S, Gard G, Dehlendorff C, Iversen HK, Biering-Soerensen F, Jensen R. Can pelvic floor muscle training improve quality of life in men with mild to moderate post-stroke and lower urinary tract symptoms? *Eur J Phys Rehabil Med*. 2017;53:416-425.
227. Tibaek S, Gard G, Jensen R. Pelvic floor muscle training is effective in women with urinary incontinence after stroke: a randomised, controlled and blinded study. *Neurourol Urodyn*. 2005;24:348-357.
228. Stoniute A, Madhuvrata P, Still M, Barron-Millar E, Nabi G, Omar MI. Oral anticholinergic drugs versus placebo or no treatment for managing overactive bladder syndrome in adults. *Cochrane Database Syst Rev*. 2023;5:Cd003781.
229. Harari D, Norton C, Lockwood L, Swift C. Treatment of constipation and fecal incontinence in stroke patients: randomized controlled trial. *Stroke*. 2004;35:2549-2555.
230. Todd CL, Johnson EE, Stewart F, et al. Conservative, physical and surgical interventions for managing faecal incontinence and constipation in adults with central neurological diseases. *Cochrane Database Syst Rev*. 2024;10:Cd002115.
231. Tuong NE, Klausner AP, Hampton LJ. A review of post-stroke urinary incontinence. *Can J Urol*. 2016;23:8265-8270.
232. Fluck A, Fry CH, Affley B, et al. Sex-specific independent risk factors of urinary incontinence in acute stroke patients: A multicentre registry-based cohort study. *Neurourol Urodyn*. 2024;43:818-825.

233. Lucente G, Corral J, Rodríguez-Esparragoza L, et al. Current Incidence and Risk Factors of Fecal Incontinence After Acute Stroke Affecting Functionally Independent People. *Front Neurol*. 2021;12:755432.

#### **Additional References for Performance Measures**

- Smith A, Hewitt J, Quinn TJ, Robling M. Patient-reported outcome measures (PROMs) use in post-stroke patient care and clinical practice: a realist synthesis protocol. *Syst Rev*. 2021 Apr 28;10(1):128. doi: 10.1186/s13643-021-01682-w. PMID: 33910631; PMCID: PMC8082773.
- Schmidt, R., Geisler, D., Urban, D. et al. Stroke survivors' preferences on assessing patient-reported outcome measures. *J Patient Rep Outcomes* 7, 124 (2023). <https://doi.org/10.1186/s41687-023-00660-1>
- Ibrahim S, Francis T, Sheehan KA, Kokorelias K, Stanimirovic A, Hashmi S, Kalocsai C, Ng S, Berkhout SG, Cameron JI, Rac V and Pikula A (2024) Exploring unmet needs and preferences of young adult stroke patients for post-stroke care through *PROMs and gender differences*. *Front. Stroke* 3:1386300. doi: 10.3389/fstro.2024.1386300