



CANADIAN STROKE BEST PRACTICE RECOMMENDATIONS

**Rehabilitation, Recovery and Community Participation
following Stroke**

**Part Three: Optimizing Activity and Community Participation
following Stroke**

7th Edition, 2025

Appendix Three: References

*Salbach N, Yao J (Writing Group Co-Chairs)
Nelson M, Shi J (Section One Co-Leads)
on behalf of the Canadian Stroke Best Practice Recommendations
Stroke Rehabilitation and Recovery Writing Group*

© 2025 Heart and Stroke Foundation of Canada

APPENDIX THREE: REFERENCES

1. Nelson MLA, Hanna E, Hall S, Calvert M. What makes stroke rehabilitation patients complex? Clinician perspectives and the role of discharge pressure. *J Comorb.* 2016;6:35-41.
2. Holodinsky JK, Lindsay P, Yu AXY, Ganesh A, Joundi RA, Hill MD. Estimating the Number of Hospital or Emergency Department Presentations for Stroke in Canada. *Can J Neurol Sci.* 2023;50:820-825.
3. Government of Canada. Canadian Chronic Disease Surveillance System (CCDSS). 2023; <https://health-infobase.canada.ca/ccdss/data-tool/Index>. Accessed March 3, 2025.
4. Kapral MK, Hall R, Fang J, et al. Predictors of Hospitalization in Patients With Transient Ischemic Attack or Minor Ischemic Stroke. *Can J Neurol Sci.* 2016;43:523-528.
5. Canadian Institutes of Health Research. Transitions in Care: Overview. 2019; <http://www.cihr-irsc.gc.ca/e/50972.html> Accessed October 31, 2024.
6. Li S, Francisco GE, Rymer WZ. A New Definition of Poststroke Spasticity and the Interference of Spasticity With Motor Recovery From Acute to Chronic Stages. *Neurorehabil Neural Repair.* 2021;35:601-610.
7. Robinson RG, Jorge RE. Post-Stroke Depression: A Review. *Am J Psychiatry.* 2016;173:221-231.
8. Taylor WD, Steffens DC, MacFall JR, et al. White matter hyperintensity progression and late-life depression outcomes. *Arch Gen Psychiatry.* 2003;60:1090-1096.
9. Chen L, Xiong S, Liu Y, et al. C-Reactive Protein Can Be an Early Predictor of Poststroke Apathy in Acute Ischemic Stroke Patients. *J Stroke Cerebrovasc Dis.* 2018;27:1861-1869.
10. Miller DS, Robert P, Ereshefsky L, et al. Diagnostic criteria for apathy in neurocognitive disorders. *Alzheimers Dement.* 2021;17:1892-1904.
11. Marin RS. Apathy: a neuropsychiatric syndrome. *J Neuropsychiatry Clin Neurosci.* 1991;3:243-254.
12. Steffens DC, Fahed M, Manning KJ, Wang L. The neurobiology of apathy in depression and neurocognitive impairment in older adults: a review of epidemiological, clinical, neuropsychological and biological research. *Transl Psychiatry.* 2022;12:525.
13. van Dalen JW, Moll van Charante EP, Nederkoorn PJ, van Gool WA, Richard E. Poststroke apathy. *Stroke.* 2013;44:851-860.
14. Lapchak PA. Neuronal Dysregulation in Stroke-Associated Pseudobulbar Affect (PBA): Diagnostic Scales and Current Treatment Options. *J Neurol Neurophysiol.* 2015;6.
15. Buysse DJ. Sleep health: can we define it? Does it matter? *Sleep.* 2014;37:9-17.

16. Acciarresi M, Bogousslavsky J, Paciaroni M. Post-stroke fatigue: epidemiology, clinical characteristics and treatment. *Eur Neurol*. 2014;72:255-261.
17. Staub F, Bogousslavsky J. Fatigue after stroke: a major but neglected issue. *Cerebrovasc Dis*. 2001;12:75-81.
18. Annoni JM, Staub F, Bogousslavsky J, Brioschi A. Frequency, characterisation and therapies of fatigue after stroke. *Neurol Sci*. 2008;29 Suppl 2:S244-246.
19. Lerdal A, Bakken LN, Kouwenhoven SE, et al. Poststroke fatigue--a review. *J Pain Symptom Manage*. 2009;38:928-949.
20. World Health Organization. International Classification of Functioning, Disability and Health. 2001; <https://iris.who.int/bitstream/handle/10665/42407/9241545429-eng.pdf>. Accessed June 12, 2024.
21. Canadian Stroke Best Practices. Canadian Stroke Best Practice Recommendations. Overview of Methodology 7th Edition 2019-2023; <https://www.strokebestpractices.ca/recommendations/overview-methods-and-knowledge-translation>. Accessed March 3, 2025.
22. Brouwers MC, Kho ME, Browman GP, et al. AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ*. 2010;182:E839-842.
23. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64:383-394.
24. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924-926.
25. Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schünemann HJ. What is "quality of evidence" and why is it important to clinicians? *BMJ*. 2008;336:995-998.
26. Schünemann H BJ, Guyatt G, Oxman A, editors. GRADE handbook for grading quality of evidence and strength of recommendations. 2013; <https://guidelinedevelopment.org/handbook>. Accessed March 3, 2025.
27. Lam RW, Kennedy SH, Adams C, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) 2023 Update on Clinical Guidelines for Management of Major Depressive Disorder in Adults: Réseau canadien pour les traitements de l'humeur et de l'anxiété (CANMAT) 2023 : Mise à jour des lignes directrices cliniques pour la prise en charge du trouble dépressif majeur chez les adultes. *Can J Psychiatry*. 2024;69:641-687.
28. Jørgensen TS, Wium-Andersen IK, Wium-Andersen MK, et al. Incidence of Depression After Stroke, and Associated Risk Factors and Mortality Outcomes, in a Large Cohort of Danish Patients. *JAMA Psychiatry*. 2016;73:1032-1040.
29. Horne KS, Gibson EC, Byrne J, Bender JR, Robinson GA. Post-stroke apathy: A case series investigation of neuropsychological and lesion characteristics. *Neuropsychologia*. 2022;171:108244.

30. Nelsone L, Rafsten L, Abzhandadze T, Sunnerhagen KS. A cohort study on anxiety and perceived recovery 3 and 12 months after mild to moderate stroke. *Front Neurol*. 2023;14:1273864.
31. Hackett ML, Pickles K. Part I: frequency of depression after stroke: an updated systematic review and meta-analysis of observational studies. *Int J Stroke*. 2014;9:1017-1025.
32. Ayerbe L, Ayis S, Crichton S, Wolfe CD, Rudd AG. The natural history of depression up to 15 years after stroke: the South London Stroke Register. *Stroke*. 2013;44:1105-1110.
33. Salinas J, Ray RM, Nassir R, et al. Factors Associated With New-Onset Depression Following Ischemic Stroke: The Women's Health Initiative. *J Am Heart Assoc*. 2017;6:e003828.
34. Guiraud V, Gallarda T, Calvet D, et al. Depression predictors within six months of ischemic stroke: The DEPRESS Study. *Int J Stroke*. 2016;11:519-525.
35. Shi Y, Yang D, Zeng Y, Wu W. Risk Factors for Post-stroke Depression: A Meta-analysis. *Front Aging Neurosci*. 2017;9:218.
36. Swartz RH, Cayley ML, Lanctôt KL, et al. The "DOC" screen: Feasible and valid screening for depression, Obstructive Sleep Apnea (OSA) and cognitive impairment in stroke prevention clinics. *PLoS One*. 2017;12:e0174451.
37. Karamchandani RR, Vahidy F, Bajgur S, et al. Early depression screening is feasible in hospitalized stroke patients. *PLoS One*. 2015;10:e0128246.
38. Meader N, Moe-Byrne T, Llewellyn A, Mitchell AJ. Screening for poststroke major depression: a meta-analysis of diagnostic validity studies. *J Neurol Neurosurg Psychiatry*. 2014;85:198-206.
39. Prisnie JC, Fiest KM, Coutts SB, et al. Validating screening tools for depression in stroke and transient ischemic attack patients. *Int J Psychiatry Med*. 2016;51:262-277.
40. Allida SM, Hsieh CF, Cox KL, et al. Pharmacological, non-invasive brain stimulation and psychological interventions, and their combination, for treating depression after stroke. *Cochrane Database Syst Rev*. 2023;7:Cd003437.
41. Xu XM, Zou DZ, Shen LY, et al. Efficacy and feasibility of antidepressant treatment in patients with post-stroke depression. *Medicine (Baltimore)*. 2016;95:e5349.
42. Allida S, Cox KL, Hsieh CF, House A, Hackett ML. Pharmacological, psychological and non-invasive brain stimulation interventions for preventing depression after stroke. *Cochrane Database Syst Rev*. 2020;5:Cd003689.
43. Yi ZM, Liu F, Zhai SD. Fluoxetine for the prophylaxis of poststroke depression in patients with stroke: a meta-analysis. *Int J Clin Pract*. 2010;64:1310-1317.
44. Robinson RG, Jorge RE, Moser DJ, et al. Escitalopram and problem-solving therapy for prevention of poststroke depression: a randomized controlled trial. *JAMA*. 2008;299:2391-2400.

45. Mikami K, Jorge RE, Moser DJ, et al. Increased frequency of first-episode poststroke depression after discontinuation of escitalopram. *Stroke*. 2011;42:3281-3283.
46. Robinson RG, Jorge RE, Long J. Prevention of Poststroke Mortality Using Problem-Solving Therapy or Escitalopram. *Am J Geriatr Psychiatry*. 2017;25:512-519.
47. Knapp P, Dunn-Roberts A, Sahib N, et al. Frequency of anxiety after stroke: An updated systematic review and meta-analysis of observational studies. *Int J Stroke*. 2020;15:244-255.
48. Knapp P, Campbell Burton CA, Holmes J, et al. Interventions for treating anxiety after stroke. *Cochrane Database Syst Rev*. 2017;5:Cd008860.
49. Chun HY, Newman R, Whiteley WN, Dennis M, Mead GE, Carson AJ. A systematic review of anxiety interventions in stroke and acquired brain injury: Efficacy and trial design. *J Psychosom Res*. 2018;104:65-75.
50. Caeiro L, Ferro JM, Costa J. Apathy Secondary to Stroke: A Systematic Review and Meta-Analysis. *Cerebrovasc Dis*. 2013;35:23-39.
51. Tan S, Lin X, Liu Z, Wu Y, Xie B, Wang T. Non-pharmacological intervention effects on apathy caused by central nervous system organic diseases: A network meta-analysis. *Medicine (Baltimore)*. 2022;101:e30467.
52. Allida S, House A, Hackett ML. Pharmaceutical interventions for emotionalism after stroke. *Cochrane Database Syst Rev*. 2022;11:Cd003690.
53. Lee EJ, Kim JS, Chang DI, et al. Depressive Symptoms in Stroke Patients: Are There Sex Differences? *Cerebrovasc Dis*. 2020;49:19-25.
54. Brown DL, Jiang X, Li C, et al. Sleep apnea screening is uncommon after stroke. *Sleep Med*. 2019;59:90-93.
55. Zhan J, Zhang P, Wen H, et al. Global prevalence estimates of poststroke fatigue: A systematic review and meta-analysis. *Int J Stroke*. 2023;18:1040-1050.
56. Cumming TB, Packer M, Kramer SF, English C. The prevalence of fatigue after stroke: A systematic review and meta-analysis. *Int J Stroke*. 2016;11:968-977.
57. Snaphaan L, van der Werf S, de Leeuw FE. Time course and risk factors of post-stroke fatigue: a prospective cohort study. *Eur J Neurol*. 2011;18:611-617.
58. Duncan F, Wu S, Mead GE. Frequency and natural history of fatigue after stroke: a systematic review of longitudinal studies. *J Psychosom Res*. 2012;73:18-27.
59. Wu S, Kutlubaev MA, Chun HY, et al. Interventions for post-stroke fatigue. *Cochrane Database Syst Rev*. 2015;2015:Cd007030.

60. Bivard A, Lillicrap T, Krishnamurthy V, et al. MIDAS (Modafinil in Debilitating Fatigue After Stroke): A Randomized, Double-Blind, Placebo-Controlled, Cross-Over Trial. *Stroke*. 2017;48:1293-1298.
61. Poulsen MB, Damgaard B, Zerahn B, Overgaard K, Rasmussen RS. Modafinil May Alleviate Poststroke Fatigue: A Randomized, Placebo-Controlled, Double-Blinded Trial. *Stroke*. 2015;46:3470-3477.
62. Choi-Kwon S, Choi J, Kwon SU, Kang DW, Kim JS. Fluoxetine is not effective in the treatment of post-stroke fatigue: a double-blind, placebo-controlled study. *Cerebrovasc Dis*. 2007;23:103-108.
63. Nguyen S, Wong D, McKay A, et al. Cognitive behavioural therapy for post-stroke fatigue and sleep disturbance: a pilot randomised controlled trial with blind assessment. *Neuropsychol Rehabil*. 2019;29:723-738.
64. Zedlitz AM, Rietveld TC, Geurts AC, Fasotti L. Cognitive and graded activity training can alleviate persistent fatigue after stroke: a randomized, controlled trial. *Stroke*. 2012;43:1046-1051.
65. Johansson B, Bjuhr H, Rönnbäck L. Mindfulness-based stress reduction (MBSR) improves long-term mental fatigue after stroke or traumatic brain injury. *Brain Inj*. 2012;26:1621-1628.
66. Seiler A, Camilo M, Korostovtseva L, et al. Prevalence of sleep-disordered breathing after stroke and TIA: A meta-analysis. *Neurology*. 2019;92:e648-e654.
67. Swartz RH, Longman RS, Lindsay MP, et al. Canadian Stroke Best Practice Recommendations: Vascular cognitive impairment, 7th edition practice guidelines update, 2024. *Alzheimers Dement*. 2025;21:e14324.
68. Sexton E, McLoughlin A, Williams DJ, et al. Systematic review and meta-analysis of the prevalence of cognitive impairment no dementia in the first year post-stroke. *Eur Stroke J*. 2019;4:160-171.
69. Cicerone KD, Langenbahn DM, Braden C, et al. Evidence-based cognitive rehabilitation: updated review of the literature from 2003 through 2008. *Arch Phys Med Rehabil*. 2011;92:519-530.
70. O'Donoghue M, Leahy S, Boland P, Galvin R, McManus J, Hayes S. Rehabilitation of Cognitive Deficits Poststroke: Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Stroke*. 2022;53:1700-1710.
71. Rogers JM, Foord R, Stolwyk RJ, Wong D, Wilson PH. General and Domain-Specific Effectiveness of Cognitive Remediation after Stroke: Systematic Literature Review and Meta-Analysis. *Neuropsychol Rev*. 2018;28:285-309.
72. Rozental-Iluz C, Zeilig G, Weingarden H, Rand D. Improving executive function deficits by playing interactive video-games: secondary analysis of a randomized controlled trial for individuals with chronic stroke. *Eur J Phys Rehabil Med*. 2016;52:508-515.
73. Chung CS, Pollock A, Campbell T, Durward BR, Hagen S. Cognitive rehabilitation for executive dysfunction in adults with stroke or other adult non-progressive acquired brain damage. *Cochrane Database Syst Rev*. 2013;2013:Cd008391.

74. Poulin V, Korner-Bitensky N, Dawson DR, Bherer L. Efficacy of executive function interventions after stroke: a systematic review. *Top Stroke Rehabil.* 2012;19:158-171.
75. Loetscher T, Potter KJ, Wong D, das Nair R. Cognitive rehabilitation for attention deficits following stroke. *Cochrane Database Syst Rev.* 2019;2019.
76. das Nair R, Cogger H, Worthington E, Lincoln NB. Cognitive rehabilitation for memory deficits after stroke. *Cochrane Database Syst Rev.* 2016;9:Cd002293.
77. Aben L, Heijenbrok-Kal MH, van Loon EM, et al. Training memory self-efficacy in the chronic stage after stroke: a randomized controlled trial. *Neurorehabil Neural Repair.* 2013;27:110-117.
78. Aben L, Heijenbrok-Kal MH, Ponds RW, Busschbach JJ, Ribbers GM. Long-lasting effects of a new memory self-efficacy training for stroke patients: a randomized controlled trial. *Neurorehabil Neural Repair.* 2014;28:199-206.
79. Oberlin LE, Waiwood AM, Cumming TB, Marsland AL, Bernhardt J, Erickson KI. Effects of Physical Activity on Poststroke Cognitive Function: A Meta-Analysis of Randomized Controlled Trials. *Stroke.* 2017;48:3093-3100.
80. Cumming TB, Tyedin K, Churilov L, Morris ME, Bernhardt J. The effect of physical activity on cognitive function after stroke: a systematic review. *Int Psychogeriatr.* 2012;24:557-567.
81. Zhang B, Li D, Liu Y, Wang J, Xiao Q. Virtual reality for limb motor function, balance, gait, cognition and daily function of stroke patients: A systematic review and meta-analysis. *J Adv Nurs.* 2021;77:3255-3273.
82. Mingming Y, Bolun Z, Zhijian L, Yingli W, Lanshu Z. Effectiveness of computer-based training on post-stroke cognitive rehabilitation: A systematic review and meta-analysis. *Neuropsychol Rehabil.* 2022;32:481-497.
83. Nie P, Liu F, Lin S, et al. The effects of computer-assisted cognitive rehabilitation on cognitive impairment after stroke: A systematic review and meta-analysis. *J Clin Nurs.* 2022;31:1136-1148.
84. Exalto LG, Weaver NA, Kuijf HJ, et al. Sex Differences in Poststroke Cognitive Impairment: A Multicenter Study in 2343 Patients With Acute Ischemic Stroke. *Stroke.* 2023;54:2296-2303.
85. Gladstone DJ, Lindsay MP, Douketis J, et al. Canadian Stroke Best Practice Recommendations: Secondary Prevention of Stroke Update 2020. *Can J Neurol Sci.* 2022;49:315-337.
86. Lim JH, Lee HS, Song CS. Home-based rehabilitation programs on postural balance, walking, and quality of life in patients with stroke: A single-blind, randomized controlled trial. *Medicine (Baltimore).* 2021;100:e27154.
87. Pang MY, Eng JJ, Dawson AS, McKay HA, Harris JE. A community-based fitness and mobility exercise program for older adults with chronic stroke: a randomized, controlled trial. *J Am Geriatr Soc.* 2005;53:1667-1674.

88. Eng JJ, Chu KS, Kim CM, Dawson AS, Carswell A, Hepburn KE. A community-based group exercise program for persons with chronic stroke. *Med Sci Sports Exerc.* 2003;35:1271-1278.
89. Saunders DH, Sanderson M, Hayes S, et al. Physical fitness training for stroke patients. *Cochrane Database Syst Rev.* 2020;3:Cd003316.
90. Shepperd S, Wee B, Straus SE. Hospital at home: home-based end of life care. *Cochrane Database Syst Rev.* 2011:Cd009231.
91. Malhotra C, Shafiq M, Batcagan-Abueg APM. What is the evidence for efficacy of advance care planning in improving patient outcomes? A systematic review of randomised controlled trials. *BMJ Open.* 2022;12:e060201.
92. Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ.* 2010;340:c1345.
93. Kirchhoff KT, Hammes BJ, Kehl KA, Briggs LA, Brown RL. Effect of a disease-specific advance care planning intervention on end-of-life care. *J Am Geriatr Soc.* 2012;60:946-950.
94. Yu S, Muhunthan J, Lindley R, et al. Driving in stroke survivors aged 18-65 years: The Psychosocial Outcomes In Stroke (POISE) Cohort Study. *Int J Stroke.* 2016;11:799-806.
95. Barco PP, Wallendorf MJ, Snellgrove CA, Ott BR, Carr DB. Predicting road test performance in drivers with stroke. *Am J Occup Ther.* 2014;68:221-229.
96. Devos H, Akinwuntan AE, Nieuwboer A, Truijen S, Tant M, De Weerd W. Screening for fitness to drive after stroke: a systematic review and meta-analysis. *Neurology.* 2011;76:747-756.
97. George S, Crotty M, Gelinas I, Devos H. Rehabilitation for improving automobile driving after stroke. *Cochrane Database Syst Rev.* 2014;2014:Cd008357.
98. Edwards JD, Kapoor A, Linkewich E, Swartz RH. Return to work after young stroke: A systematic review. *Int J Stroke.* 2018;13:243-256.
99. Hackett ML, Glozier N, Jan S, Lindley R. Returning to paid employment after stroke: the Psychosocial Outcomes In Stroke (POISE) cohort study. *PLoS One.* 2012;7:e41795.
100. Hannerz H, Holbæk Pedersen B, Poulsen OM, Humle F, Andersen LL. A nationwide prospective cohort study on return to gainful occupation after stroke in Denmark 1996-2006. *BMJ Open.* 2011;1:e000180.
101. Trygged S, Ahacic K, Kåreholt I. Income and education as predictors of return to working life among younger stroke patients. *BMC Public Health.* 2011;11:742.
102. Cain S, Churilov L, Collier JM, et al. Factors associated with paid employment 12 months after stroke in A Very Early Rehabilitation Trial (AVERT). *Ann Phys Rehabil Med.* 2022;65:101565.

103. Wang YC, Kapellusch J, Garg A. Important factors influencing the return to work after stroke. *Work*. 2014;47:553-559.
104. Ntsiea MV, Van Aswegen H, Lord S, Olorunju SS. The effect of a workplace intervention programme on return to work after stroke: a randomised controlled trial. *Clin Rehabil*. 2015;29:663-673.
105. Baldwin C, Brusco NK. The effect of vocational rehabilitation on return-to-work rates post stroke: a systematic review. *Top Stroke Rehabil*. 2011;18:562-572.
106. Brouns R, Valenzuela Espinoza A, Goudman L, Moens M, Verlooy J. Interventions to promote work participation after ischaemic stroke: A systematic review. *Clin Neurol Neurosurg*. 2019;185:105458.
107. Orange C, Lanhers C, Coll G, et al. Determinants of Return to Work After a Stroke: A Systematic Review and Meta-analysis. *Arch Phys Med Rehabil*. 2024;105:359-368.
108. Jee S, Sohn MK, Lee J, et al. Prediction for return to driving after the first-ever stroke in Korea: The KOSCO study. *J Rehabil Med*. 2018;50:800-805.
109. Lee D, Heffron JL, Mirza M. Content and Effectiveness of Interventions Focusing on Community Participation Poststroke: A Systematic Review. *Arch Phys Med Rehabil*. 2019;100:2179-2192.e2171.
110. Dorstyn D, Roberts R, Kneebone I, Kennedy P, Lieu C. Systematic review of leisure therapy and its effectiveness in managing functional outcomes in stroke rehabilitation. *Top Stroke Rehabil*. 2014;21:40-51.
111. Harrington R, Taylor G, Hollinghurst S, Reed M, Kay H, Wood VA. A community-based exercise and education scheme for stroke survivors: a randomized controlled trial and economic evaluation. *Clin Rehabil*. 2010;24:3-15.
112. Desrosiers J, Noreau L, Rochette A, et al. Effect of a home leisure education program after stroke: a randomized controlled trial. *Arch Phys Med Rehabil*. 2007;88:1095-1100.
113. Stein J, Hillinger M, Clancy C, Bishop L. Sexuality after stroke: patient counseling preferences. *Disabil Rehabil*. 2013;35:1842-1847.
114. Buzzelli S, di Francesco L, Giaquinto S, Nolf G. Psychological and medical aspects of sexuality following stroke. *Sex Disabil*. 1997;15:261-270.
115. Sansom J, Ng L, Zhang N, Khan F, Couldrick L. Let's talk about sex: A pilot randomised controlled trial of a structured sexual rehabilitation programme in an Australian stroke cohort. *International Journal of Therapy and Rehabilitation*. 2015;22:21-29.
116. Guo M, Bosnyak S, Bontempo T, et al. Let's Talk About Sex! - Improving sexual health for patients in stroke rehabilitation. *BMJ Qual Improv Rep*. 2015;4.
117. Song H, Oh H, Kim H, Seo W. Effects of a sexual rehabilitation intervention program on stroke patients and their spouses. *NeuroRehabilitation*. 2011;28:143-150.

118. Stratton H, Sansom J, Brown-Major A, Anderson P, Ng L. Interventions for sexual dysfunction following stroke. *Cochrane Database Syst Rev*. 2020;5:Cd011189.
119. Gall SL, Tran PL, Martin K, Blizzard L, Srikanth V. Sex differences in long-term outcomes after stroke: functional outcomes, handicap, and quality of life. *Stroke*. 2012;43:1982-1987.
120. Chen C, Reeves MJ, Lisabeth LD. Sex Differences in Participation Restriction in Social Activities Among Older Stroke Survivors: A Nationwide Study. *Stroke*. 2025;56:265-275.

Additional References for Performance Measures

- Smith A, Hewitt J, Quinn TJ, Robling M. Patient-reported outcome measures (PROMs) use in post-stroke patient care and clinical practice: a realist synthesis protocol. *Syst Rev*. 2021 Apr 28;10(1):128. doi: 10.1186/s13643-021-01682-w. PMID: 33910631; PMCID: PMC8082773.
- Schmidt, R., Geisler, D., Urban, D. et al. Stroke survivors' preferences on assessing patient-reported outcome measures. *J Patient Rep Outcomes* 7, 124 (2023).
<https://doi.org/10.1186/s41687-023-00660-1>
- Ibrahim S, Francis T, Sheehan KA, Kokorelias K, Stanimirovic A, Hashmi S, Kalocsai C, Ng S, Berkhout SG, Cameron JI, Rac V and Pikula A (2024) Exploring unmet needs and preferences of young adult stroke patients for post-stroke care through *PROMs and gender differences*. *Front. Stroke* 3:1386300. doi: 10.3389/fstro.2024.1386300