



CANADIAN STROKE BEST PRACTICE RECOMMENDATIONS


Stroke Systems of Care **7th Edition, Update 2026** **Evidence Tables** ***Stroke Recognition and Response***

*Sacha Arsenault and Gary Hunter (Writing Group Co-Chairs)
on Behalf of the Canadian Stroke Best Practice Recommendations
Stroke Systems of Care Writing Group*

Table of Contents

Search Strategy	3
Published Guidelines	4
Awareness of Stroke Signs & Symptoms among the General Public	6
Increasing Awareness of Stroke Symptoms through Public Health Campaigns	11
References.....	25

Search Strategy



PubMed, EMBASE and the Cochrane Central Register of Controlled Trials databases were search using terms such as “stroke risk factors”, “stroke symptoms”, “stroke awareness”, “stroke behaviours”, “stroke campaigns”, “public education” and “awareness campaigns”. Titles and abstract of each article were reviewed for relevance. Bibliographies were reviewed to find additional relevant articles. Articles were excluded if they were: non-English, commentaries, case-studies, narrative, book chapters, editorials, non-systematic review, or conference abstracts. Additional searches for relevant best practice guidelines were completed and included in a separate section of the review. A total of 32 articles and 5 guidelines were included and were separated into separate categories designed to answer specific questions.

Published Guidelines

Guideline	Recommendations
<p>National Clinical Guideline for Stroke for the UK and Ireland. London: Intercollegiate Stroke Working Party; 2023 May 4.</p> <p>Available at: www.strokeguideline.org.</p>	<p>Recommendation Public awareness campaigns of the symptoms of stroke should be recurrent, targeted at those most at risk of stroke, and formally evaluated. [2016]</p>
<p>Powers WJ, Rabinstein AA, Ackerson T, Adeoye OM, Bambakidis NC, Becker Ket al; on behalf of the American Heart Association Stroke Council.</p> <p>Guidelines for the early management of patients with acute ischemic stroke: 2019 Update to the 2018 Guidelines for the Early Management of Acute Ischemic Stroke: A Guideline for Healthcare Professionals from the American Heart Association/American Stroke Association</p> <p><i>Stroke</i> 2019;50:e344–e418.</p>	<ol style="list-style-type: none"> 1. Public health leaders, along with medical professionals and others, should design and implement public education programs focused on stroke systems and the need to seek emergency care (by calling 9-1-1) in a rapid manner. These programs should be sustained over time and designed to reach racially/ethnically, age, and sex diverse populations. (Class 1; LOE B-NR). 2. Such educational programs should be designed to specifically target the public, physicians, hospital personnel, and emergency medical services (EMS) personnel to increase use of the 9-1-1 EMS system, to decrease stroke onset to emergency department (ED) arrival times, and to increase timely use of thrombolysis and thrombectomy. (Class I: LOE C-EO). 3. Activation of the 9-1-1 system by patients or other members of the public is strongly recommended. 9-1-1 dispatchers should make stroke a priority dispatch, and transport times should be minimized. (Class 1; LOE B-NR).
<p>Adeoye O, Nystrom KV, Yavagal DR, et al.</p> <p>Recommendations for the Establishment of Stroke Systems of Care: A 2019 Update.</p> <p><i>Stroke</i> 2019; 50: e187-e210.</p>	<ol style="list-style-type: none"> 1. Public health leaders along with medical professionals and others should design and implement public education programs focused on stroke systems and the need to seek emergency care (by calling 9-1-1) in a rapid manner. These programs should be repetitive and designed to reach diverse populations. Further research is needed to establish the most effective programs for diverse populations. (New) 2. EMS leaders, in coordination with local, regional, and state agencies and in consultation with medical authorities and local experts, should develop triage paradigms and protocols that ensure that all patients with a known or suspected stroke are rapidly identified and assessed with a validated and standardized instrument for stroke screening such as FAST (Face, Arm, Speech, Time), Los Angeles Prehospital Stroke Screen, or CPSS. (Revised) 3. When there are several intravenous alteplase-capable hospitals in a well-defined geographic region, extra transportation times to reach a facility capable of endovascular thrombectomy should be limited to no more than 15 minutes in patients with a prehospital stroke severity scale score suggestive of LVO. When several hospital options exist within similar travel times, EMS should seek care at the facility capable of offering the highest level of stroke care. Further research is needed to establish travel time parameters for hospital bypass in cases of prehospital suspicion of LVO. (New)

Guideline	Recommendations
<p>Kobayashi A, Czlonkowska A, Ford GA, Fonseca AC, Luijckx GJ, Korv J, et al.</p> <p>European Academy of Neurology - European Stroke Organisation consensus statement and practical guidance for pre-hospital management of stroke.</p> <p><i>Eur J Neurol</i> 2018 Mar;25(3):425-433.</p>	<p>We recommend educational campaigns to increase the awareness of immediately calling EMS for people with suspected stroke. (SOR strong; QOE very low quality of evidence)</p>
<p>Clinical Guidelines for Stroke Management 2017. Melbourne (Australia): National Stroke Foundation.</p>	<p>Not included in update</p>

Evidence Tables

Awareness of Stroke Signs & Symptoms among the General Public

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
Dharmasaroja & Uransilp 2024 Thailand Cross-sectional survey	NA	281 participants ≥40 years, without a history of stroke. Mean age was 58 years, 77% were women. 60% had ≥ 1 vascular risk factor.	Data were gathered from two electronic surveys, which were divided into three parts: i) symptoms of acute stroke based on BE FAST, with 5 possible correct answers (B [balance: gait ataxia/leg weakness], E [eyes: visual symptoms], F [facial droop], A [arm weakness], S [speech disturbance], ii) vascular risk factors (7 correct answers; hypertension, diabetes, hyperlipidemia, smoking, alcohol drinking, heart diseases), and iii) how they should respond (time to come to ER after stroke onset and emergency calling number).	Primary outcome: Knowledge	133 participants (47%) had good knowledge of stroke awareness and were able to identify 3-5 signs of stroke. The percentage of respondents who correctly identified signs of stroke Balance problems: 117 (42%) Eyes: 110 (39%) Facial droop: 177 (63%) Arm weakness: 181 (64%) Speech: 153 (54%) Independent predictors of having good knowledge of stroke symptoms were age > 50 years, higher education and female sex. For the question about how fast they should come to the ER if they suspected an acute stroke Immediately: 182 (65%) Within 4.5 hours: 43 (15%) Within 6 hours: 3 (1%), within 24 hours: 18 (6%) Within 3 days: 1 (0.4%) Did not know: 34 (12%) Respondents who had good knowledge of stroke symptoms also had good knowledge of vascular risk factors and knew to reach the Emergency Department immediately.

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
<p>Naguib et al. 2020 Saudi Arabia Cross sectional study Awareness about Stroke and Proper Actions to Be Taken; A room for Improvement</p>	NA	600 adults recruited using convenience sampling at public shopping malls from different areas in Riyadh, from September 2018 until March 2019. Mean age was 27.7 years, 39.5% were women. 50.2% had attended university.	Participants were interviewed to determine whether they could recognize ≥ 5 stroke risk factors, 3 warning signs of stroke, and whether they could call an ambulance in the first 3 hours of a suspected incident. Those that fulfilled all criteria were considered knowledgeable.	Primary outcome: Knowledge	<p>40% of respondents were considered knowledgeable. Independent factors associated with being knowledgeable were younger age, female, and being employed.</p> <p>The majority of respondents correctly recognized hypertension (64.7%), heart diseases (61%), smoking (58.2%), dyslipidemia (57.7%), increasing age (55.2%), and obesity (50%) as stroke risk factors.</p> <p>68.3% of participants were able to identify at least 3 warning signs of stroke including slurred speech (61.8%), sudden weakness in arms, legs or face (54.5%), and sudden tingling or loss of sensation in 1 side of the body (53.8%). 9.5% of respondents were not able to identify any warning signs.</p> <p>70% of respondents indicated they would call an ambulance within 3 hours of observed symptoms, while 86.2% indicated they would call a physician within 24 hours.</p>
<p>Krishnamurthi et al. 2020 New Zealand Survey</p>	NA	400 adults, aged ≥ 20 years in New Zealand, stratified by the 4 main ethnic groups (Māori, Pacific, Asian/ other, and Europeans), selected through landline telephone directory listings. 42% of participants were between 40 and 59 years of age, had completed at least high school education (97%), were employed (69%), and lived in urban areas (92%)	The survey was conducted between April and June 2015 by telephone interview. Participants were asked to answer a series of questions on stroke knowledge and awareness, using unprompted and prompted methods of questioning	Primary outcome: Knowledge, predictors of greater knowledge	<p>Response rate was 57%.</p> <p>Using a prompted method, sudden speech difficulty (94%) and sudden weakness on 1 side (92%) were the most commonly identified stroke symptoms.</p> <p>When asked “Do you think you would be able to tell if someone was having a stroke?” 57% answered yes, while 43% did not believe they could.</p> <p>When prompted with a list of 11 options, 98% of participants identified 1 stroke symptom, 96% identified 2 or more, 86% identified 3 or more, 64% identified 4 or more and 37% identified all 5 of the correct symptoms in the list.</p> <p>Without prompting, 78% of participants correctly identified ≥ 1 risk factor, 62% identified ≥ 2, and 35% identified ≥ 3.</p>

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
					Higher education level, higher income, and personal experience of stroke were predictive of greater awareness.
Metias et al. 2017 Canada Survey	NA	Adult patients, accompanying family and friends who were in the waiting room of community- and academic-based vascular clinics in Toronto, at 2 time periods 2009-2010 and 2014-2015	Questionnaires were administered to participants, including questions pertained to knowledge of stroke risk factors and symptoms. Participants were also asked to identify their response to an individual who appeared to be experiencing a stroke, stroke treatments and stroke resources that were available. The responses to these questions were used to develop a stroke literacy score. 'Stroke literacy' was established by listing ≥1 stroke risk factor and ≥1 signs of stroke.	Primary outcome: Stroke literacy	In 2010, 207 surveys were distributed, of which 198 were completed. The mean age of participants was 60.5 years. In 2015, 791 questionnaires were completed. The mean age of participants was 53.4 years. The most frequently identified risk factors for stroke were smoking in 2010 and hypertension in 2015. The most common stroke symptom identified was trouble speaking (56.6%) in 2010 and weakness, numbness or paralysis (67.1%) in 2015. 80.3% vs. 83.1% of participants were able to identify ≥1 stroke risk factor in 2010 and 2015, respectively. The corresponding values for ability to identify a stroke sign/symptom were 90.9% and 88.7%. Television was the most common source of information. The Heart & Stroke Foundation of Canada was a source of information for 6.6% of participants in 2010 compared with 35.8% of those in 2015.
Mochari-Greenberger et al. 2014 USA Cross-sectional survey	NA	1,205 women aged ≥25 years living in the United States, who had participated in the American Heart Association National Women's Tracking Survey.	Participants were contacted by telephone through random-digit dialing and asked standardized questions, related to stroke warning signs, actions to take in the event of stroke.	Primary outcome: Knowledge	The percentage of women who recognized the following signs of stroke: Sudden weakness/numbness of the face or limb of one side: 51%. Loss of/trouble with understanding speech:44% Sudden severe headache: 23% Unexplained dizziness: 20% Loss of vision in one eye: 18%

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
					One in 5 women could not identify one stroke warning sign.
Miyamatsu et al. 2013 Japan Cross-sectional survey	NA	5,540 participants, aged 40-74 years, randomly selected from the Basic Resident Register, recruited from 3 large cities in Japan	A multiple choice, mail-in survey including items related to general knowledge of stroke, early symptoms of stroke, information sources and what to do if a stroke is suspected.	Primary outcome: Knowledge	<p>The response rate was 49.0%.</p> <p>Mean age was 58 years. 53% of participants were female.</p> <p>Recognition of stroke symptoms:</p> <p>Sudden one-sided weakness: 86.6%</p> <p>Sudden confusion: 86.6%</p> <p>Sudden headache: 72.3%</p> <p>Sudden dizziness: 62.7%</p> <p>Sudden trouble seeing: 35.0%</p> <p>All 5 symptoms correctly identified: 23.0%</p> <p>81.2% of participants indicated they would call an ambulance immediately in response to a suspected stroke.</p> <p>Mass media campaigns (particularly those appearing on television) were identified as the most common source of information (estimated fraction: 0.32), followed by newspapers (estimated fraction: 0.28). The internet and personal communication were the least common sources of information.</p>
Robinson et al. 2013 UK Cross-sectional survey	NA	1,300 individuals (39% male) representative of the general population were sampled from public areas, places of work, and academic institutions.	Items on the survey represented the following: 1) basic familiarity with the concept of stroke, 2) awareness of warning signs of stroke, 3) awareness of risk factors for stroke, and 4) knowledge of the FAST campaign.	Primary outcome: Awareness of stroke warning signs and risk factors and knowledge of the FAST campaign.	<p>70% of those surveyed were aware of the FAST campaign and 80% recalled the 'burning face' image.</p> <p>Over 75% of participants were able to recall all three FAST stroke symptoms and >90% were able to recall at least one.</p>

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
					<p>Stroke warning signs not included as part of the FAST campaign were not as well recognized (e.g., Visual loss=44%, Dizziness=47%).</p> <p>Hypertension, smoking, alcohol, and diabetes were identified as risk factors for stroke by 90%, 74%, 54%, and 51% of participants, while 68% of participants were aware that stroke affects the brain.</p>
<p>Lundelin et al. 2012</p> <p>Spain</p> <p>Cross-sectional survey</p>	NA	<p>11,827 non-institutionalized adults living in Spain who had participated in the Study on Nutrition & Cardiovascular Risk in Spain (ENRICA) study</p>	<p>Participants were identified by multistage clustered random sampling. Data was collected using a telephone interview. 55% of the participants were also examined physically and provided a blood sample.</p>	<p>Primary outcome: Stroke knowledge and predictors of knowledge, based on sociodemographic indicators</p>	<p>65.2% of participants could correctly identify 4-6 symptoms of stroke. 19% could identify all 6 symptoms correctly, while 11.4% were unable to identify a single symptom.</p> <p>Higher levels of education were associated with better knowledge.</p> <p>81.1% of participants indicated that they would call an ambulance if they suspected someone was having a stroke. Persons who could identify more stroke symptoms were more likely to call for an ambulance.</p>
<p>Hickey et al. 2009</p> <p>Ireland</p> <p>Cross-sectional survey</p>	NA	<p>2033 community-dwelling older adults. 57% female.</p> <p>The survey response rate was 68%.</p>	<p>Participants completed a survey with items addressing knowledge of stroke warning signs and risk factors.</p>	<p>Primary outcome: Percentage of participants correctly identifying stroke warning signs and risk factors.</p>	<p>Warning signs: Slurred speech (54%), dizziness (44%), numbness (41%), weakness (38%), headache (29%), and vision problems (20%).</p> <p>Risk Factors: hypertension (75%), cholesterol (40%), smoking (30%), diabetes (11%), and alcohol use (10%).</p>

Increasing Awareness of Stroke Symptoms through Public Health Campaigns

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
Darkhabani et al. 2024 USA RCT	CA: <input checked="" type="checkbox"/> Blinding: Patient <input checked="" type="checkbox"/> Assessor <input checked="" type="checkbox"/> ITT: <input checked="" type="checkbox"/>	174 participants recruited from the general public through community-based events using study-specific flyers advertisements and brochures distributed at a booth during events. Median age was 56 years, 25% were men. 43% were in the medical profession, 55% had a family member with a history of stroke, 90% were able to list stroke symptoms prior to any education.	Participants were randomized to an educational session on either FAST or BE FAST using printed cards representing each letter of the mnemonic.	Primary outcome: Mnemonic recall at 3-5 minutes, 60 minutes and 30 days	134 (77%) participants completed the 30-day assessment. There were no significant differences between groups in mnemonic recall between FAST and BE FAST education groups at any of the time points (99.4% at 3-5 minutes, 89.1% at 60 minutes and 74.7% at 30 days). A significantly higher percentage of participants recalled all items in the FAST group (75% vs. 30.2% at 3-5 minutes; 70.5% vs. 41.9% 60 minutes and 51.1% vs. 24.4% at 30 days). Retention was significantly higher for some of the individual items of FAST at some of the assessment points. Mnemonic recall of FAST and BE FAST at 3-5 minutes, 60 minutes and 30 days was similar between participants with a medical and nonmedical background.
Rioux et al. 2022 Canada Cross sectional surveys	NA	2,451 unique respondents from 4 waves of surveys conducted between July 2016 and January 2019 in Quebec. Most people were aged ≥35 years. 51.5% were women.	Knowledge of FAST stroke signs was assessed at each wave administered by telephone, which were preceded by public awareness campaigns. A bilingual English/French FAST public awareness campaign preceded survey waves 1-3 and two campaigns preceded wave 4. Participants were asked open-ended questions	Primary outcomes: Stroke symptom recognition and intention to call EM	There was a significant decrease in the percentage of persons who could not name a single FAST sign, from 37.7% in wave 1, to 26.9% in wave 2, to 30.4% in wave 3 to 30.5% in wave 4 (p<0.001). Speech difficulty was the most frequently reported sign (54.4%), followed by arm weakness (53.8%) and face drooping in the last survey (36.2%). In the last survey, 2 factors were independently associated with worse performance on FAST identification: male sex and retired (vs. working). There was an increase over time in the percentages of participants who were familiar with the FAST campaign (34.5%, 45.1%, 53.2% and 49.3%, p<0.001).

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
			on stroke definition, their first action in the event of a stroke, the main signs that a person is having a stroke and whether they recalled the preceding FAST campaign.		There was an increase over time in the percentages of participants who were able to provide a correct definition of stroke (54.3%, 66.1%, 71.1% and 67.0%, p<0.001). The percentage of participants indicating they would call EMS/911 as a first response to a suspected stroke did not change significantly over time (81.5%, 80.9%, 85.5% and 80.1%, p=0.158).
Tan et al. 2022 Singapore Systematic review & meta-analysis	All studies were assessed as being at high risk of bias.	13 studies including 113,592 adults aged < 65 years recruited from the general population.	The effects of stroke education using mass media campaigns on stroke symptom recognition and intention to call emergency medical services, were estimated.	Primary outcome: Stroke symptom recognition and intention to call EM	Duration of the campaigns ranged from 3 weeks to two years. 4 studies used FAST as the basis of their campaigns. Mass media campaigns increased the likelihood of symptom recognition compared with pre-campaign (RR=1.20 (95% CI 1.07-1.36, n=5 studies). Mass media campaigns increased the likelihood that persons would call EMS, compared with pre-campaign (RR= 1.19, 95% CI 1.11 -1.28, 8 studies included).
Han & Jang 2021 South Korea Cochrane review (protocol)	NA	RCTs including members of the general public.	Trials will examine public education interventions using mass media to increase awareness of stroke signs and symptoms and what to do in the event of a stroke.	Primary outcome: Knowledge of stroke symptoms, knowledge of what to do in the event of stroke to reduce prehospital delays Secondary outcomes: Use of thrombolytics, stroke-related mortality, stroke severity, and satisfaction with the intervention	TBD

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
Hickey et al. 2018 UK Survey	NA	1,010 randomly selected adults from a nationally representative sample (response rate of 50.6%). 48% were men.	Data collected before a public awareness campaign in 2009 (wave 1) were compared with a post-awareness campaign in 2011 (wave 2), through telephone surveys, including the same participants from waves 1 and 2.	Primary outcomes: Knowledge (≥ 2 stroke warning signs or risk factors for stroke), intended response to witnessing a stroke	<p>Compared with wave 1, there were no significant differences in the percentage of participants who could correctly describe a stroke, who described observable signs of a stroke, who could give a partially correct description of a stroke or who didn't know what a stroke was following wave 2. A significantly lower percentage of participants in wave 2 gave an incorrect description a stroke (10% vs. 17%).</p> <p>Following wave 2, a significantly higher percentage of participants could identify correctly high blood pressure, and poor diet as risk factors for stroke (31% vs. 27% and 28% vs. 19%, respectively) with no improvement in those recognizing smoking, lack of exercise, high cholesterol, heredity, or increasing age as risk factors.</p> <p>Following wave 2, a significantly increased percentage of participants identified all the FAST components as signs of stroke, with significantly fewer participants recognizing dizziness, visual problems or severe headache as signs of stroke. The ability to identify weakness on one side of the body was increased significantly in wave 2, with no differences between waves for unilateral numbness, difficulty understanding, loss of movement, disorientation, and loss of coordination.</p> <p>Significantly fewer participants were unable to identify any signs of stroke following wave 2 (4% vs. 23%).</p> <p>Following wave 2, a significantly higher percentage of participants indicated they would call an ambulance (57.2% vs. 47.1%), but with significantly fewer indicating the need to get other medical help (14.6% vs. 24.2%).</p> <p>In both waves, 1 and 2, men were less likely than women to correctly identify ≥ 2 signs of stroke.</p>

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
					Respondents with a previous stroke or myocardial infarction were more likely to identify signs of stroke.
Wolters et al. 2018 UK Prospective study	NA	2,243 consecutive patients from the e OxVasc study presenting with first TIA or stroke, seeking medical attention between April 1, 2002, and March 31, 2014. Mean age was 73.6, 50.2% were women.	Response to the FAST campaign (television + public transit displays), which ran from 2009 intermittently though 2014, was assessed.	Primary outcomes: Number of early recurrent strokes in patients who delayed or failed to seek medical attention, and the odds of seeking urgent attention after TIA and minor stroke before vs after initiation of the public education campaign.	825 (36.8%) patients were initially seen with TIA, 831 (37.0%) with minor stroke, and 587 (26.2%) with major stroke. For major stroke, the use of nonemergency services declined significantly from 2009 onward, while the use of EMS increased significantly (58.8% before April 1, 2009, vs 78.9% after April 1, 2009, p=0.001). In addition, first medical attention was sought more quickly (within 3 hours) after April 1, 2009 (67.6% vs. 81.3%; OR=2.08; 95% CI, 1.40-3.11; P < .001). Time to first seeking medical attention after TIA and minor stroke was similar before and after April 1, 2009, with 42.1% seeking medical attention within 3 hours prior to April 1, 2009 vs. 40.4%, after (OR= 0.79; 95% CI, 0.50-1.23). The corresponding values for patients seeking help within 24 hours of symptoms before and after the FAST campaign were similar (70.4% vs. 70.6%; OR=0.75; 95% CI, 0.48-1.19). 185 patients had a stroke within 90 days of their initial TIA or stroke, of whom 93 (49.5%) followed unheeded TIAs for which no medical attention was sought. The number of strokes preceded by an unheeded TIA was similar before and after the FAST campaign (8.0% vs. 8.1%, p=0 .93).
Nordanstig et al. 2017 Sweden Prospective study	NA	1,500 randomly selected persons, aged 15 to 79 years from the Swedish population	The effect of a 27-month long public awareness campaign, designed to increase knowledge of the Swedish translation of FAST and, the need to call the Swedish	Primary outcomes: Knowledge of words in FAST mnemonic, need to call emergency services	The response rate decreased from 62% to 36% over time. 52% of respondents had seen as advertisement during the campaign, compared with 29% prior. From pre-campaign (survey 1) to end of the campaign (survey 8), the number of persons who:

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
			<p>emergency number, was evaluated. The campaign included television advertisements, spots on public service television, and paid advertisements and banners in newspapers and social media, plus a website and Facebook page. Telephone interviews were carried out with participants at 8 points: before, 3 times during, immediately after, and 9, 13 and 21 months after the campaign.</p>		<p>1) had heard of FAST increased from 15%-50% (adj OR=1.30, 95% CI 1.26-1.33), 2) could recall all keywords in the mnemonic increased from 0.3% to 2% (adj OR=1.34, 95% CI 1.15-1.55, p<0.0001), 3) could recall some/all keywords in the mnemonic had increased from 4% to 14% (adj OR=1.23, 95% CI 1.17-1.28, p<0.0001) and 4) intended to call emergency services increased from 65% to 73% (adj OR=1.05, 95% CI 1.03-1.08, p<0.0001).</p> <p>From during and directly after the campaign, to 21 months after the campaign, there was no change in the number of respondents who had heard about the campaign (51% vs 50%). During the same timeframe, the number of persons who: 1) could recall all keywords in the mnemonic decreased significantly from 3.4% to 2.3%, p=0.043, 2) could recall some/all keywords in the mnemonic had decreased significantly from 23% to 14%, p<0.0001, and 3) intended to call emergency services decreased significantly from 76% to 73%, p=0.037.</p>
<p>Advani et al. 2016 Norway Prospective study</p>	<p>NA</p>	<p>All patients admitted to the ER with signs and symptoms of stroke</p>	<p>The effects of a 1-month mass media intervention (MMI), using advertising on television, posters, social media and healthcare trust websites was used to promote signs of stroke symptoms, using the FAST mnemonic to encourage people to seek immediate medical attention.</p> <p>1,400 telephone surveys were also conducted,</p>	<p>Primary outcome: Changes in t-PA use after the MMI</p> <p>Secondary outcome: Changes in the numbers of stroke admissions to the ER within 4.5 hours of stroke onset</p>	<p>The average number of patients treated with t-PA increased significantly from 7.3 to 11.3 patients per month (an increase of 54.7%, p=0.02) during the period from the 12 months preceding the MMI to the 6 months after the MMI (including the month during which the campaign ran).</p> <p>After the initial 6-month period, the average number of patients treated with t-PA dropped to 9.5, which was still significantly higher than the 12 months preceding the MMI.</p> <p>The average number of patients treated in the ER increased significantly from 37.3 to 72.8 patients per month (an increase of 95.7%, p<0.001) during the period from the 12 months preceding the MMI</p>

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
			before and after the campaign		<p>to the 6 months after the MMI (including the month during which the campaign ran).</p> <p>After the initial 6-month period, the average number of patients treated in the ER dropped to 52.6, which was still significantly higher than the 12 months preceding the MMI.</p> <p>Telephone survey results: the number of people who could name any stroke symptom increased from 66% to 75% after the MMI.</p> <p>Of those who could name a symptom, 52% recognized facial droop, 42% named speech difficulties and 42% named arm weakness.</p>
Bray et al. 2015 Australia Cross-sectional study	NA	NA	The association between monthly volumes of ambulance dispatches for stroke and 12 National Stroke Foundation multimedia regional public awareness campaigns (2004-2014) was explored. The campaigns lasted on average for 6 weeks and were both paid and pro bono. Most of the campaigns focused on FAST symptoms and the need for quick response	Primary outcome: Changes in ambulance calls for stroke following public awareness campaigns, controlling for paid funding, timing of campaign exposure, number of days in the month, season, population size and population growth	<p>11/12 campaigns were associated with increases in call volumes. The percentage increases by year were:</p> <p>2004: 4.8%</p> <p>2005: 7.6%</p> <p>2006: -2.2%</p> <p>2007: 1.0%</p> <p>2008: 5.2%</p> <p>2009: 4.7%</p> <p>2010: 6.9%</p> <p>2011: 5.7%</p> <p>2014: 9.9% (campaign 1); 9.3% (campaign 2)</p> <p>The effect persisted for approximately 3 months.</p> <p>In 2014, one of the campaigns was run nationally. There was an increase of 1478 calls (6.7%).</p> <p>The campaigns run in urban areas tended to be more successful than those in rural settings.</p>

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
					Compared with regions that did not receive funding, those that did reported a relative increase of 10.2% in call volumes.
Wolters et al. 2015 UK Prospective study	NA	688 participants in the OXVASC study who had experienced a major stroke (NIHSS>3). Mean age was 77 years, 57% were female. Median stroke severity was 9.	Patient behavior for out of hospital strokes was compared before (2002-2008) and after (2009-2013) the introduction of UK-FAST (a public education television campaign Feb-Apr 2009)	Primary outcome: Time from stroke onset to seeking medical attention and the type of medical attention sought.	There were 416 strokes pre-FAST and 252 post-FAST. Medical attention was sought by a by-stander in the majority of cases (553, 89.6%) The median time to seek first medical attention decreased significantly post-FAST (53 vs. 31 minutes, p=0.005). Median time to arrival to hospital decreased significantly post-FAST (185 vs. 119 minutes, p<0.0001). First contact with EMS or ED was made more frequently post FAST (74.8% vs. 57.2%, OR=2.20, 95% CI 1.55-3.13, p<0.0001), while first contact with non-emergency services (e.g., GP) decreased significantly (42.8% vs. 25.2%, OR=0.45, 95% CI 0.32-0.65, p<0.0001). The number of patients who arrived at hospital within 3 hours increased significantly post FAST (46.9% vs. 65.8%, OR=2.18, 95% CI 1.55-3.06, p<0.0001).
Flynn et al. 2014 UK Time series evaluation	NA	Data from 3 databases (Stroke Association, Hospital Episode Statistics (HES) for England, and the Safe Implementation of Thrombolysis in Stroke UK (SITS-UK) database.	The impact of the 3 national FAST campaigns, (Feb-Mar 2009, Nov-Dec 2009 and Feb-Mar 2010) was assessed over time, using data from 3 databases	Primary outcome: Information-seeking behavior for stroke, emergency admissions, 911 calls, number of patients receiving thrombolytic treatment	There was significantly increased activity on the Stroke Associations' website following the first campaign (webpage views), information materials dispatched and calls to the help line, with decreases after the campaign ceased. The same pattern of activity was observed during campaigns 2 and 3. There were significantly greater number of emergency admissions over the 22-month period prior to the first campaign, and significant increases during the first campaign, but the increases thereafter were not significant.

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
					There was a significant increase in the number of patients treated with t-PA prior to the first campaign, with a non-significant increase following it. Following the first campaign, there was an increase of 3 patient/month treated with t-PA and again after the third campaign (3 patients/month).
Rasura et al. 2014 Italy Review	NA	22 intervention studies and 5 web-based stroke education campaigns. 14 studies targeted the general public using mass media campaigns, which varied in duration from 3 months to 4 years. 6 studies targeted specific groups with the interventions lasting 3 minutes-12 hours.	Narrative synthesis of included studies	Primary outcome: Not stated <i>a priori</i>	3 popular stroke signs and symptoms were included in all of the studies using mass media campaigns: FAST, SUDDEN and Give-Me-Five. Effectiveness of the interventions was assessed in most studies through questionnaires administered pre and post intervention. Increases of emergency room presentations t-PA administration and ambulance dispatches, and reductions in pre-hospital delays were also used. The authors concluded that large public health campaigns using mass media are expensive and short lived and may not be effective, although the increased costs can be mitigated through more prompt treatment with t-PA. Lower cost, smaller scale educational campaigns can be delivered successfully in the community. The message being delivered must direct the person to call an ambulance. The dose of the campaign appears to be as important as the message. Television appears to be the most effective medium. Online campaigns can also be successful but tend to attract a self-selected group (e.g. well-educated women).
Dombrowski et al. 2014 UK RCT	CA: ☒ Blinding: Patient ☒	5,000 adults, sampled from the electoral register in a large urban community	Participants were randomized to receive a questionnaire + an Act FAST leaflet delivered	Primary outcome: Awareness of the Act FAST campaign, and if so, what the acronym stands for.	Data from 1615 respondents (32.3%) were available. Mean age was 54 years. 57% were female.

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
	Assessor <input checked="" type="checkbox"/> ITT: <input checked="" type="checkbox"/>		by mail or to the questionnaire only. The Act FAST campaign was a mass media campaign designed to increase awareness of the FAST mnemonic	Stroke response and recognition were assessed using 16 vignette scenarios (12 stroke and 4 non-stroke)	A higher proportion of participants in the leaflet group had heard of the Act FAST campaign (75% vs. 68%, p<0.001). Significantly more persons in the leaflet group correctly named the FAST elements (66% vs. 45%, p<0.001). There was no difference between groups in the number or participants who indicated they would call "999" in the event of a stroke. Using the 16 vignettes, there were no significant differences between groups in the numbers of participants who were able to correctly identify stroke symptoms, not even in the 6 FAST scenarios (78.4% correct recognition in the leaflet group vs. 79.0% in the non-leaflet group, p=0.55). The pattern of correct response was similar (65.6% vs. 64.0%, p=0.30).
Mellon et al. 2014 UK Retrospective study	NA	870 patients who had presented to the emergency department (ED) of 2 large teaching hospitals with symptoms of stroke over a one-year time frame (March 2010-Feb 2011)	Multivariable analysis to determine the impact of 3 waves of 3-week mass media FAST campaigns in May, August 2010 and January 2011	Primary outcome: Factors related to onset of symptoms to ER arrival ≤3.5 hours and changes in presentations to the ED with stroke symptoms.	There were 284 confirmed strokes (32.6%) and 150 TIAs (17.4%). The remaining admissions were for non-stroke neurological and medical conditions. Independent predictors of arrival to ED ≤ 3.5 hours were arrival via ambulance (OR=3.1, 95% CI 2.23-4.23, p<0.001) and self-referral (OR=2.67, 95% CI 1.84-3.88, p<0.001). Exposure to the FAST campaign was not a significant predictor (OR=0.93, 95% CI 0.60-1.45, p=0.76). During the first wave of the campaign, there was an increase in the number of presentations to the ED with stroke symptoms, which dropped off after the cessation of the campaign. There was no such evidence of an impact of the FAST campaign during waves 2 or 3.
Bray et al. 2013 Australia	NA	12,439 participants, ≥40 years of age randomly selected from the	Surveys were administered via telephone over a 6-year	Primary outcome: Number of respondents aware of	From 2004 to 2010, a significant increase was observed in the number of respondents aware of the campaign (31% vs 50%) and in the number or

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
Cross-sectional survey		general population using an electronic telephone directory.	period to determine the impact of a national multimedia stroke awareness campaign on knowledge of stroke warning signs, and awareness of the campaign.	the advertising campaign, number of respondents able to recall ≥ 1 , ≥ 2 , and/or ≥ 3 stroke warning signs.	participants able to name ≥ 1 (69% vs 81%), ≥ 2 (43% vs 63%), and ≥ 3 (19% vs 32%) warning signs of stroke (all $p < 0.001$). Respondents who could identify ≥ 2 warning signs were significantly more likely to be aware of the campaign (OR= 1.88, 95% CI 1.74 to 2.04).
Trobbiani et al. 2013 Australia Cross-sectional surveys (pre/post intervention)	NA	English sample: 1905 adults aged 16-55 years Canadian sample: 2807 adults ≥ 18 years Australian sample: 1,002 adults >40 years, randomly selected from Victorian households	Comparison of 3 mass media stroke awareness campaigns conducted in England (FAST), Canada (SUDDENS) and Australia (FAST), designed to increase the public's recognition of stroke symptoms. The surveys were conducted by telephone interviews (Canada, Australia) or in person (England) before and after exposure to the campaigns. All of the campaigns used television and radio, print newspapers and digital means.	Primary outcome: Changes in the ability to recognize stroke symptoms following awareness campaigns. Timing of data collection surveys (before/after campaign): Canada: 1 month/2 months Australia: 10 months/2 weeks England: 6 weeks/1 month	The cost of the campaigns was €7 million (England), €700K (Canada) and €650K (Australia). Head-to-head comparisons across the 3 groups was not possible. Prior to the campaign, a significantly greater number of English participants could identify more FAST components compared with Australian participants (F: 67% vs. 44%, A: 62% vs. 30%, S: 56% vs. 21%, T: 42% vs. 14%, all $p < 0.001$). Following the campaign significantly more English participants had increased their knowledge (F: 19% vs. 3%, A: 12% vs. 6%, S: 10% vs. 6%, T: 6% vs. -1%, all $p < 0.01$). Prior to the campaign, a significantly greater number of Australian participants could identify ≥ 2 warning signs of stroke compared with Canadian participants (59% vs. 50%, $p < 0.001$). Following the campaign, the percent change (increase) was similar between groups (Canada 7% vs. Australia 9%). Higher percentages of participants in England and Australia indicated they would call emergency services in the event of a stroke, compared with those from Canada (97% and 90% vs. 67%).
Worthmann et al. 2013	NA	1,004 adult residents of the city of Hanover	Computer-assisted interviews were conducted before and	Primary outcome: Changes in the ability to recognize stroke	There was a significant increase in the number of participants who identified paresis and weakness as signs of stroke (from 40% to 46%, $p = 0.007$ and

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
Germany Cross-sectional surveys (pre/post intervention)			<p>immediately after a 6-month public awareness campaign, designed to increase knowledge of stroke knowledge warning signs and risk factors, emergency care seeking behavior.</p> <p>The campaign included mass media (posters, flyers, public lectures, newspapers and television advertisements).</p>	<p>symptoms following the awareness campaign.</p>	<p>24% to 31%, p<0.001, respectively). There was also a significant increase in the number of participants who identified impaired vision as a sign of stroke (8% to 11%, p<0.05).</p> <p>There were no significant increases in the number of participants who identified gait disorders, numbness, sudden headache, nausea or disorientation as a sign of stroke.</p> <p>Following the campaign, there was a significant increase in the number of participants who indicated that the first action to take after recognizing a possible stroke was to call for emergency care (from 74% to 84%, p<0.001). There was a non-significant increase in the number of participants who indicated this action should be taken immediately (from 81% to 84%)</p>
Fogle et al. 2010 USA Controlled study	NA	<p>Intervention group: 400 adults randomly selected, living in a single county with a census of 85,314. Median age was 40 years.</p> <p>Control group: 401 adults randomly selected, living in a comparable county, with a census of 81,763. Median age was 33 years.</p>	<p>Participants in the intervention group were exposed to a high-intensity public education campaign, conducted during 2, 10-week periods (2007-2008). The campaign included television, radio and newspaper advertisements that focused on stroke warning signs and actions to take.</p> <p>Participants in the control group were not exposed to the campaign.</p> <p>Computer-assisted telephone surveys of</p>	<p>Primary outcome:</p> <p>Changes in the ability to recall ≥2 stroke warning signs before and after campaign, stroke risk factors and intentions to call '911' if a stroke was suspected</p>	<p>There was a significant increase in the number of participants in the intervention group, but not the control group, who could identify ≥2 warning signs before/after the campaign (from 73% to 82% and 68% to 69%, respectively).</p> <p>There was a significant increase in the number of participants in the intervention group, but not the control group, who could identify ≥2 risk factors for stroke before/after the campaign (from 69% to 86% and 70% to 65%, respectively).</p> <p>There was a non-significant increase in the number of participants in the intervention group who indicated that they would call '911' if they suspected they, or someone else were having a stroke (from 81% to 84%), while there was a significant decrease among participants in the control group (from 82% to 74%, p<0.05).</p>

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
			adults ≥45 years were conducted before and after the campaign.		
Lecoutuier et al. 2010 UK Systematic Review	NA	10 studies examining the effectiveness of mass media campaigns designed to change stroke-related knowledge and/or behaviour following the onset of stroke. 4 studies targeted the public and 4 targeted both the public and healthcare professionals (HCP).	Narrative synthesis	Primary outcome: Knowledge of stroke symptoms and the need for rapid response, access to emergency services, and early treatment with thrombolysis.	Each of the 6 studies targeting public awareness demonstrated a significant increase in knowledge of stroke symptoms; however, these interventions appeared to have little impact on emergency response behaviours. Four studies targeted both professionals and the public. In light of the dual purpose of these interventions, the authors concluded that it was difficult to “disentangle any active components that might explain any reported impact”. The interventions appeared to be more effective for HCP than the public.
Jurkowski et al. 2010 USA Controlled study	NA	Intervention group: 994 adults ≥30 years, randomly selected, living in 3 counties with a total population of 603K. Mean age was 56 years. 65% female Control group: 795 adults living in one county with a total population of 376K. Mean age was 54 years. 63% female.	Participants in the intervention group were exposed to a 3-phase multimedia campaign, designed to increase public awareness of the FAST mnemonic. Each stage of the campaign lasted for 33 weeks Participants in the control group were not exposed to the campaign. Telephone surveys were conducted before and during phase 3 of the campaign.	Primary outcome: Percentage of respondents aware of the campaign, aware of the campaigns primary message, and who would call 9-1-1 in response to specific stroke symptoms identified in oneself and/or others.	The percentage of respondents who reported they would call 9-1-1 in response to specific stroke symptoms increased significantly more from pre-to post campaign for participants in the intervention group, with increases ranging from 9%-12% for specific symptoms identified in oneself and 4%-12% for specific symptoms identified in others.
Marx et al. 2010 Germany	NA	501 adults residing in 5 counties in western Germany. The census	Computer-assisted telephone surveys were conducted before and after a 3-month mass-	Primary outcome: Gender differences in general stroke knowledge,	Prior to the intervention, significantly more women than men could correctly answer the question “where does stroke happen in the body?” (87.2% vs. 70.3%, p<0.001) and knew the stroke

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
Surveys (pre/post intervention)		from the area was 400K. Mean age was 52 years. 44% male.	media educational, program (billboard, flyers, radio, television, and 3 full day public events) designed to increase awareness of stroke signs and symptoms and response.	knowledge of stroke warning signs and risk factors, response in the event of stroke and through what source(s) had participants gained this knowledge	<p>emergency call number (33.3% vs. 24.4%, p<0.001). Significantly more women knew to call for emergency care (87.2% vs. 70.1%, p<0.001)</p> <p>Following the intervention, an increased number of both men and women could answer the 2 questions correctly. The percentage change from pre to post intervention was significantly higher for women, (Question 1: +2.8%, p=n/s; Question 2, +5.7%, p<0.005).</p> <p>There were increases in the mean number of stroke warning signs that could be named before and after the intervention (women: 5.4 to 6.2; men: 5.1 to 5.9).</p> <p>Men and women were equally likely to recall campaign messages from television, radio and flyers. Women were more likely to remember the messages from leaflets and advertisements in pharmacies and doctor's offices, while men were more likely to remember them from advertisements on buses and street cars.</p>
Hodgson et al. 2007 Canada Retrospective study	NA	1,000 adults ≥45 years residing in the province of Ontario.	Telephone surveys were conducted before, during and following 2 paid television advertising campaigns that lasted for 8 and 9 months, conducted in 2003 and 2005, designed to increase awareness of the warning signs of stroke.	Primary outcome: Identification of up to 5 stroke warning signs, and the proportion of participants who could name ≥2 signs of stroke	<p>Over the 6 sampling points, there was a significant increase in the proportion of participants who could correctly identify ≥2 stroke warning signs, and the mean number of warning signs (p<0.001).</p> <p>Prior to the first campaign, 52.1% of participant could correctly identify ≥2 stroke warning signs. Following the completion of the first campaign, this percentage increased to 67.8%.</p> <p>From part-way through the second campaign to 7 months following its completion, there was a significant decrease in the percentage of persons who could correctly identify ≥2 stroke warning signs (70.8% to 64.2%, p<0.001).</p> <p>Across all polls, significantly more women than men could name ≥2 stroke warning signs.</p>

Study/Type	Quality Rating	Sample Description	Method	Outcomes	Key Findings and Recommendations
					Only a small percentage of participants could identify all 5 signs of stroke (paralysis, slurred speech, headache, blurred vision and dizziness). (1.3%, prior to the first campaign, to a maximum of 3.5%, immediately following the second campaign)
Morgenstern et al. 2007 USA RCT Kids Identifying and Defeating Stroke (KIDS)	CA: ☒ Blinding: Patient ☒ Assessor ☒ ITT: ☒	294 students in 3 intervention schools and 279 students in 3 control schools. There were significantly more males in the intervention schools (55% vs. 41%, p=0.002)	Students in the intervention group received 4, 50-minute classroom-based lessons for 3 years, starting in grades 6, designed to increase stroke awareness and response. Homework assignments were designed to include (and educate) the student's parents. Students in the control group received no stroke education.	Primary outcome: 3 domains (4 questions each): stroke pathophysiology (domain 1), stroke symptom knowledge (domain 2) and behavioral intent to call 911 in the event of stroke (domain 3).	47% of students in the intervention group and 46% in the control group completed the pre and post tests. Only 18% and 16% of the parents completed both tests. (no analyses were conducted). There was a significant increase in the proportion of students answering questions correctly in domains 2 and 3 in both groups; however, the improvement was significantly greater for students in the intervention group. For domain 1, there was an increase in correct responses from 29% to 34% in the intervention group and a significant decrease in the control group (from 28% to 25%, p=0.007).

Abbreviations

BE FAST: Balance, Eye, Face, Arm, Speech, Time	CA: concealed allocation	CI: confidence interval
EMS: Emergency Medical Services	FAST: face, arm, speech, time	HTN: hypertension
ITT: intention-to-treat	RR: relative risk	OR: odds ratio

References

- Advani R, Naess H, Kurz M. Mass media intervention in western Norway aimed at improving public recognition of stroke, emergency response, and acute treatment. *J Stroke Cerebrovasc Dis* 2016;25(6):1467-72.
- Bray JE, Johnson R, Trobbiani K, Mosley I, Lalor E, Cadilhac D. Australian public awareness of stroke warning signs improves after national multimedia campaigns. *Stroke* 2013;44:3540-3543.
- Bray JE, Straney L, Barger B, Finn J. Effect of public awareness campaigns on calls to ambulance across Australia. *Stroke* 2015;46(5):1377-1380.
- Darkhabani MZ, Homa-Bonell JK, Thoreson L, Bobholz JA, Spaulding D, Engebose M. BE FAST Versus FAST: A Randomized Pilot Trial Comparing Retention of Stroke Symptoms Between 2 Mnemonics. *J Am Heart Assoc.* 2024 Oct;13(19):e035696.
- Dharmasaroja P, Uransilp N. Stroke Awareness and Knowledge in the At-Risk Population: A Community-Based Study. *Cureus.* 2024 Apr 7;16(4):e57756.
- Dombrowski SU, White M, Mackintosh JE. The stroke 'Act FAST' campaign: Remembered but not understood? *Int J Stroke* 2015; 10(3):324-30.
- Flynn D, Ford GA, Rodgers H, Price C, Steen N, Thomson RG. A time series evaluation of the FAST National Stroke Awareness Campaign in England. *PLoS One* 2014;9(8):e104289.
- Fogle CC, Oser CS, McNamara MJ, et al. Impact of media on community awareness of stroke warning signs: a comparison study. *J Stroke Cerebrovasc Dis* 2010;19:370-75.
- Han JM, Jang B-H. Educational interventions for promoting stroke literacy in the general public (Protocol). *Cochrane Database of Systematic Reviews* 2021, Issue 9. Art. No.: CD014640
- Hickey A, O'Hanlon A, McGee H, Donnellan C, Shelley E, Horgan F, O'Neill D. Stroke awareness in the general population: knowledge of stroke risk factors and warning signs in older adults. *BMC Geriatrics* 2009; 9(1): A35.
- Hickey A, Mellon L, Williams D, Shelley E, Conroy RM. Does stroke health promotion increase awareness of appropriate behavioural response? Impact of the face, arm, speech and time (FAST) campaign on population knowledge of stroke risk factors, warning signs and emergency response. *Eur Stroke J.* 2018 Jun;3(2):117-125.
- Hodgson C, Lindsay P, Rubini F. Can mass media influence emergency department visits for stroke? *Stroke* 2007;38:2115-22.
- Jurkowski, JM, Maniccia, DM, Spicer DA, Dennison BA. Impact of a multimedia campaign to increase intention to call 9-1-1 for stroke symptoms, upstate New York 2006-2007. *Preventing Chronic Disease*, 2010;7(2):A35.
- Krishnamurthi RV, Barker-Collo S, Barber PA, Tippet LJ, Dalrymple-Alford JC, Tunnage B et al. Community knowledge and awareness of stroke in New Zealand. *J Stroke Cerebrovasc Dis* 2020 Mar;29(3):104589.

- Lecouturier J, Rodgers H, Murtagh MJ, White M, Ford GA, Thomson RG. Systematic review of mass media interventions designed to improve public recognition of stroke symptoms, emergency response, and early treatment. *BMC Public Health* 2010;10:784-794.
- Lundelin K, Graciani A, Garcia-Puig J, et al. Knowledge of stroke warning symptoms and intended action in response to stroke in Spain: a nationwide population-based study. *Cerebrovasc Dis* 2012;34:161-68.
- Marx JJ, Klawitter B, Faldum A, et al. Gender-specific differences in stroke knowledge, stroke risk perception and the effects of an educational multimedia campaign. *J Neurol* 2010;257:367-74.
- Mellon L, Hickey A, Doyle F et al. Can a media campaign change health service use in a population with stroke symptoms? Examination of the first Irish stroke awareness campaign. *Emerg Med J* 2014;31:536-540.
- Metias MM, Eisenberg N, Clemente MD, Wooster EM, Dueck AD, Wooster DL, Roche-Nagle G. Public health campaigns and their effect on stroke knowledge in a high-risk urban population: A five-year study. *Vascular* 2017; 25: 497-503.
- Miyamatsu N, Okamura T, Nakayama H, et al. Public awareness of early symptoms of stroke and information sources about stroke among the general Japanese population: the Acquisition of Stroke Knowledge Study. *Cerebrovasc Dis* 2013;35:241-49.
- Mochari-Greenberger H, Towfighi A, Mosca L. National women's knowledge of stroke warning signs, overall and by race/ethnic group. *Stroke* 2014;45:1180-82.
- Morgenstern LB, Gonzales NR, Maddox KE, et al. A randomized, controlled trial to teach middle school children to recognize stroke and call 911: The kids identifying and defeating stroke project. *Stroke* 2007;38:2972-78.
- Naguib R, Fayed A, AlFadhliah AB, AlMansour NS, AlDakheel RM, AlQahtani RM. Awareness about stroke and proper actions to be taken; a room for improvement. *J Stroke Cerebrovasc Dis*. 2020 Jun;29(6):104794.
- Nordanstig A, Asplund K, Norrving B, Wahlgren N, Wester P, Rosengren L. Impact of the Swedish National Stroke Campaign on stroke awareness. *Acta Neurol Scand* 2017;136(4):345-51.
- Rasura M, Baldereschi M, Di CA, et al. Effectiveness of public stroke educational interventions: a review. *Eur J Neurol* 2014;21:11-20.
- Rioux B, Brissette V, Marin FF, Lindsay P, Keezer MR, Poppe AY. The Impact of Stroke Public Awareness Campaigns Differs Between Sociodemographic Groups. *Can J Neurol Sci*. 2022 Mar;49(2):231-238.
- Robinson TG, Reid A, Haunton VJ, Wilson A, Naylor AR. The face arm speech test: does it encourage rapid recognition of important stroke warning symptoms. *Emerg Med J* 2013;30:467-471.
- Tan J, Ramazanu S, Liaw SY, Chua WL. Effectiveness of Public Education Campaigns for Stroke Symptom Recognition and Response in Non-Elderly Adults: A Systematic Review and Meta-Analysis. *J Stroke Cerebrovasc Dis*. 2021 Nov 26;31(2):106207.

- Trobbiani K, Freeman K, Arango M, Lalor E, Jenkinson D, Thrift AG. Comparison of stroke warning sign campaigns in Australia, England, and Canada. *Int J Stroke* 2013;8 Suppl A100:28-31.
- Wolters FJ, Paul NL, Li L, Rothwell PM. Sustained impact of UK FAST-test public education on response to stroke: a population-based time-series study. *Int J Stroke* 2015. Oct;10(7):1108-14.
- Wolters FJ, Li L, Gutnikov SA, Mehta Z, Rothwell PM. Medical attention seeking after transient ischemic attack and minor stroke before and after the UK Face, Arm, Speech, Time (FAST) Public Education Campaign: Results from the Oxford Vascular Study. *JAMA Neurol* 2018; 75: 1225-1233.
- Worthmann H, Schwartz A, Heidenreich F, et al. Educational campaign on stroke in an urban population in Northern Germany: Influence on public stroke awareness and knowledge. *Int J Stroke* 2013;8:286-92.